



Atmospheric oxalic acid and related secondary organic aerosols in Qinghai Lake, a continental background site in Tibet Plateau



Jingjing Meng^a, Gehui Wang^{a,b,*}, Jianjun Li^a, Chunlei Cheng^a, Junji Cao^a

^aState Key Laboratory of Loess and Quaternary Geology, Institute of Earth Environment, Chinese Academy of Sciences, Xi'an 710075, China

^bSchool of Human Settlements and Civil Engineering, Xi'an Jiaotong University, Xi'an 710049, China

HIGHLIGHTS

- Unique molecular characteristics of SOA in remote continental atmosphere.
- Origins and formation mechanisms of SOA in Tibet Plateau.
- Biomass combustion process produces low levels of oxalate, glyoxal and methylglyoxal.

ARTICLE INFO

Article history:

Received 26 April 2013

Received in revised form

10 July 2013

Accepted 12 July 2013

Keywords:

Oxalic acid

Secondary organic aerosols

PM_{2.5}

Qinghai Lake

Tibet Plateau

ABSTRACT

Summertime PM_{2.5} aerosols collected from Qinghai Lake (3200 m a.s.l.), a remote continental site in the northeastern part of Tibetan Plateau, were analyzed for dicarboxylic acids (C₂–C₁₁), ketocarboxylic acids and α -dicarbonyls. Oxalic acid (C₂) is the dominant dicarboxylic acid in the samples, followed by malonic, succinic and azelaic acids. Total dicarboxylic acids ($231 \pm 119 \text{ ng m}^{-3}$), ketocarboxylic acids ($8.4 \pm 4.3 \text{ ng m}^{-3}$), and α -dicarbonyls ($2.7 \pm 2.1 \text{ ng m}^{-3}$) at the Tibetan background site are 2–5 times less than those detected in lowland areas such as 14 Chinese megacities. Compared to those in other urban and marine areas enhancements in relative abundances of C₂/total diacids and diacids-C/WSOC of the PM_{2.5} samples suggest that organic aerosols in the region are more oxidized due to strong solar radiation. Molecular compositions and air mass trajectories demonstrate that the above secondary organic aerosols in the Qinghai Lake atmosphere are largely derived from long-range transport. Ratios of oxalic acid, glyoxal and methylglyoxal to levoglucosan in PM_{2.5} aerosols emitted from household burning of yak dung, a major energy source for Tibetan in the region, are 30–400 times lower than those in the ambient air, which further indicates that primary emission from biomass burning is a negligible source of atmospheric oxalic acid and α -dicarbonyls at this background site.

© 2013 Elsevier Ltd. All rights reserved.

1. Introduction

Water-soluble organic compounds are important components of tropospheric aerosols, accounting for up to 70% of the total aerosol mass (Jacobson et al., 2000; Sorooshian et al., 2007a; Jung et al., 2010). Water-soluble organic compounds can change the hygroscopic properties of particles and thus influence the atmospheric radiative forcing through acting as cloud condensation nuclei (CCN). Moreover, water-soluble organic compounds such as alcohols and carboxylic acids can act as surfactants, and thus increase the solubility of airborne particulate pollutants in human respiratory tract

(Latif and Peter, 2004). Dicarboxylic acids, ketocarboxylic acids and α -dicarbonyls are important classes of water-soluble organic compounds, which are taken as secondary organic aerosols (SOA) because these compounds in the atmosphere are mostly produced from photochemical oxidation. Dicarboxylic acids and related compounds have been found in a wide variety of environments including urban (Wang et al., 2002, 2006b, 2012; Kawamura and Yasui, 2005), suburban (He and Kawamura, 2010; Ho et al., 2010), mountain (Wang et al., 2009; Hegde and Kawamura, 2012), marine (Wang et al., 2006c; Mochida et al., 2003) and free troposphere (Sorooshian et al., 2007b). Most studies have focused on dicarboxylic acids in urban atmosphere. However, information on dicarboxylic acids in high altitude regions is very limited, where dicarboxylic acids and related water-soluble species are easily lofted into clouds and activated as CCN. Tibet Plateau is called as “the roof of the world”, where population is sparse and anthropogenic

* Corresponding author. State Key Laboratory of Loess and Quaternary Geology, Institute of Earth Environment, Chinese Academy of Sciences, Xi'an 710075, China. Tel.: +86 29 8832 9320; fax: +86 29 8832 0456.

E-mail addresses: wanggh@ieecas.cn, gehuiwang@yahoo.com.cn (G. Wang).

activity is insignificant. Our previous study on atmospheric aerosols from Qinghai Lake, a remote site in northeast Tibet Plateau, found that atmospheric environment of Qinghai Lake is still pristine (Li et al., 2013). To the best of our knowledge, the current study is the first time to characterize secondary organic aerosols (SOA) in the Tibetan atmosphere. We first explored molecular composition and concentration of airborne particulate dicarboxylic acids and related compounds and then discussed their sources and formation mechanisms by comparing with those in other regions and in biomass burning emissions.

2. Experimental section

2.1. Collection of ambient aerosol

Qinghai Lake, the largest closed interior saline lake in China, is located in the northeastern part of Tibetan Plateau with an altitude of 3200 m (Fig. 1). PM_{2.5} samples were collected on the rooftop (~20 m above ground level) of a tower at the “Bird Island” peninsula (location: 36°58′44″N, 99°54′24″E), which is situated on the northwestern shore of Qinghai Lake (shown in Fig. 1). PM_{2.5} samples were collected for two months each lasting for 24 h from the beginning of July to the end of August 2010 using a high-volume air sampler (Anderson, USA) equipped with prebaked (450 °C, 8 h) quartz fiber filters (Whatman, USA) at an airflow rate of 1.13 m³ min⁻¹. Field blank samples were also collected for about 10 min without sucking any air before and after sampling by mounting a blank filter onto the sampler. After sampling, each sample was sealed in an aluminum foil bag and stored at -18 °C before analysis. A total of 34 samples collected every other day were analyzed for dicarboxylic acids and related SOA, and the remaining were reserved for a carbon isotope analysis in the future.

2.2. Collection of fresh aerosol from yak dung burning

Yak dung is the major energy source for Tibetan in the region. To investigate the impact of biomass burning emission on dicarboxylic acids and related compounds in the Qinghai Lake atmosphere, fresh PM_{2.5} aerosols emitted from household yak dung burning were

collected using the high-volume sampler above. Three local Tibetan families were randomly selected and the PM_{2.5} sampler was fixed 1 m downwind of the exhaust outlet of the domestic chimney. Each samples was collected for about 15 min and stored together with the above ambient samplers at -18 °C prior to analysis.

2.3. Sample analysis

2.3.1. Dicarboxylic acids, ketocarboxylic acids, and α -dicarbonyls

Dicarboxylic acids, ketocarboxylic acids and α -dicarbonyls were determined using the method described by Kawamura and Sakaguchi (1999). Briefly, aliquot of the filter was cut in pieces and extracted with pure Milli-Q water under ultrasonication for three times each in 15 min. The water extracts were subsequently filtrated with quartz wool and concentrated to dryness, followed by a reaction with 14% BF₃/n-butanol at 100 °C for 1 h to form butyl esters/dibutoxy acetals. After the reaction the derivatives were dissolved in n-hexane and washed with pure water for three times. Finally, the hexane layer was concentrated and analyzed using a capillary gas chromatography (GC; HP 6890) equipped with a split/splitless injector and a flame ionization detector. The GC oven temperature was programmed from 50 (2 min) to 120 °C at 15 °C min⁻¹ and then to 300 at 5 °C min⁻¹ with a final isothermal hold at 300 °C for 16 min. Peak identification was performed by comparing the GC retention time with that of authentic standards and confirmed by mass spectrum of the sample using a GC–mass spectrometry (GC–MS).

Recovery experiment was performed by spiking the authentic standards onto a blank filter and treated same as the real samples. The recoveries were 75% for oxalic and higher than 85% for other target compounds. The target compounds in the field blanks were less than 4% of those in the ambient samples. Concentrations of all determined compounds reported here were corrected for both field blanks and recoveries.

2.3.2. Inorganic ions, elemental carbon (EC), organic carbon (OC) and water-soluble organic carbon (WSOC)

Analysis details for inorganic ions, EC, OC and WSOC have been reported in our previous paper (Li et al., 2013). Briefly, aliquot of the




Fig. 1. A map description for the location of the sampling site (Bird Island) and the 72-h backward trajectories of air masses arrival at Qinghai Lake during the sampling period (local time, 24 h interval).

PM_{2.5} samples were extracted with Milli-Q water under ultrasonication for three times, and filtered through PTFE filters, then measured using an ion chromatography (Dionex 600, Dionex, USA). EC and OC in the samples were measured following the IMPROVE thermal reflectance (TOR) protocol using a DRI model 2001 carbon analyzer. As for WSOC in the samples aliquot of the filter was also extracted with pure water and analyzed using Shimadzu TOC-L CPH Total Carbon Analyzer. Data about the inorganic ions, EC, OC, WSOC and levoglucosan have been reported by Li et al. (2013) and cited here to reveal the sources and formation mechanisms of dicarboxylic acids and related secondary organic aerosols (SOA) in the region.

3. Results and discussion

3.1. Molecular characteristics of dicarboxylic acids and related SOA

A series of dicarboxylic acids and related SOA including dicarboxylic acids (C₂–C₁₁), ketocarboxylic acids and α -dicarbonyls in the PM_{2.5} samples were determined. Concentrations of these

organic compounds were summarized in Table 1, along with WSOC, EC and OC. Molecular compositions of these SOA are shown in Fig. 2.

As seen in Table 1 total dicarboxylic acids in the Qinghai Lake PM_{2.5} samples ranged from 73 to 573 ng m⁻³ with an average of 231 ± 119 ng m⁻³, which is about 2 times lower than those in mountain regions such as Mount Tai, east China (day: 756 ± 139 ng m⁻³; night: 263 ± 123 ng m⁻³) (Wang et al., 2009) and the central Himalayan in Nainital, India (Hegde and Kawamura, 2012) (day: 480 ± 132 ng m⁻³, night : 380 ± 107 ng m⁻³) and 5 times lower than in 14 Chinese megacities (892 ± 457 ng m⁻³ (Ho et al., 2007)) but around 4 times higher than those in the Pacific Ocean areas (e.g. west Pacific Ocean, 60 ng m⁻³ (Wang et al., 2006c); western North Pacific Ocean, 54 ng m⁻³ (Mochida et al., 2003)), indicating that on a global scale atmospheric SOA is much more abundant in a continental background area than in marine regions. Same as those in the above regions, the dominant species in the Qinghai Lake samples is oxalic acid (C₂), whose concentration is 139 ± 82 ng m⁻³ (29–348 ng m⁻³). Concentrations (14–139 ng m⁻³, Table S2) of C₂–C₄ in the Qinghai Lake atmosphere

Table 1
Concentrations of dicarboxylic acids, ketocarboxylic acids, α -dicarbonyls, WSOC, OC and EC in PM_{2.5} aerosols collected at Qinghai Lake, Tibet Plateau during the summer of 2010.

Compounds	Chemical formula	Concentration		
		Total (n = 34)	Easterly (n = 25)	Westerly (n = 9)
I. Dicarboxylic acids (ng m⁻³)				
Oxalic, C ₂	HOOC–COOH	139 ± 82 (29–348)	154 ± 86 (29–348)	98 ± 52 (49–199)
Malonic, C ₃	HOOC–CH ₂ –COOH	25 ± 15 (9.3–78)	28 ± 16 (9.3–78)	17 ± 7.1 (10–31)
Succinic, C ₄	HOOC–(CH ₂) ₂ –COOH	134 ± 7.8 (3.5–29)	15 ± 8.2 (3.6–29)	11 ± 5.8 (3.5–21)
Glutaric, C ₅	HOOC–(CH ₂) ₃ –COOH	2.7 ± 1.9 (0.4–6.6)	3.0 ± 2.0 (0.5–6.6)	1.9 ± 1.4 (0.4–4.6)
Adipic, C ₆	HOOC–(CH ₂) ₄ –COOH	1.1 ± 0.9 (0.2–4.9)	1.2 ± 1.0 (0.2–4.9)	1.0 ± 0.4 (0.4–1.8)
Pimelic, C ₇	HOOC–(CH ₂) ₅ –COOH	1.2 ± 0.5 (0.2–2.8)	1.2 ± 0.5 (0.2–2.8)	1.4 ± 0.3 (0.9–1.8)
Suberic, C ₈	HOOC–(CH ₂) ₆ –COOH	0.3 ± 0.2 (0.1–1.4)	0.3 ± 0.3 (0.1–1.4)	0.3 ± 0.2 (0.2–0.6)
Azelaic, C ₉	HOOC–(CH ₂) ₇ –COOH	10 ± 4.0 (5.0–22)	10 ± 4.2 (5.0–22)	11 ± 3.3 (5.6–5.6)
Sebacic, C ₁₀	HOOC–(CH ₂) ₈ –COOH	1.6 ± 0.9 (5.7–5.1)	1.6 ± 1.0 (0.7–5.1)	1.4 ± 0.3 (1.0–1.9)
Undecanedioic, C ₁₁	HOOC–(CH ₂) ₉ –COOH	2.8 ± 2.0 (1.6–13)	2.8 ± 2.3 (1.6–13)	2.9 ± 0.8 (1.6–3.7)
Methylmalonic, iC ₄	HOOC–CH(CH ₃)–COOH	1.4 ± 2.1 (0.1–12)	1.6 ± 2.4 (0.1–12)	0.8 ± 0.7 (0.1–2.4)
Methylsuccinic, iC ₅	HOOC–CH(CH ₃)–CH ₂ –COOH	0.9 ± 0.5 (0.2–2.4)	1.0 ± 0.5 (0.3–2.4)	0.7 ± 0.5 (0.2–1.4)
Methylglutaric, iC ₆	HOOC–CH(CH ₃)–(CH ₂) ₂ –COOH	0.7 ± 0.5 (0.1–2.6)	0.8 ± 0.5 (0.1–2.6)	0.6 ± 0.3 (0.2–1.1)
Maleic, M	HOOC–CH = CH–COOH(cis)	4.2 ± 2.0 (1.1–10)	4.4 ± 2.1 (1.1–10)	3.8 ± 1.5 (1.3–5.6)
Fumaric, F	HOOC–CH = CH–COOH(trans)	3.8 ± 3.0 (0.2–14)	4.2 ± 3.3 (0.2–14)	2.7 ± 1.7 (0.5–6.2)
Methylmaleic, mM	HOOC–C(CH ₃) = CH–COOH(cis)	4.0 ± 3.3 (1.3–21)	4.1 ± 3.8 (1.3–21)	3.7 ± 1.0 (2.4–5.6)
Phthalic, Ph	HOOC–C ₆ H ₄ –COOH(ortho)	6.2 ± 3.6 (2.6–22)	6.2 ± 4.1 (2.6–22)	6.3 ± 1.8 (3.4–8.0)
Isophthalic, iPh	HOOC–C ₆ H ₄ –COOH(meta)	1.4 ± 0.7 (0.8–4.4)	1.4 ± 0.8 (0.8–4.4)	1.5 ± 0.4 (0.9–2.1)
Terephthalic, tPh	HOOC–C ₆ H ₄ –COOH(para)	1.8 ± 1.1 (0.8–5.3)	1.7 ± 1.0 (0.8–5.1)	2.1 ± 1.3 (0.9–5.3)
Ketomalonic, kC ₃	HOOC–HC(O)–COOH	4.4 ± 1.4 (3.0–7.2)	4.2 ± 1.3 (3.0–7.2)	5.2 ± 1.4 (3.1–5.6)
Ketopimelic, kC ₇	HOOC(CH ₂) ₂ HC(O)(CH ₂) ₂ COOH	4.4 ± 1.7 (2.9–9.4)	4.0 ± 1.3 (2.9–7.9)	5.5 ± 2.1 (2.9–9.4)
Subtotal		231 ± 119 (73–573)	250 ± 127 (73–573)	178 ± 75 (109–331)
II. Ketocarboxylic acids (ng m⁻³)				
Pyruvic, Pyr	CH ₃ –(O)C–COOH	1.0 ± 0.5 (0.2–2.2)	1.0 ± 0.5 (0.2–2.2)	0.9 ± 0.6 (0.4–1.9)
Glyoxylic, ω C ₂	OHC–COOH	4.9 ± 3.1 (0.2–11)	5.4 ± 3.3 (0.2–13)	3.4 ± 2.1 (1.3–7.3)
3-Oxopropanoic, ω C ₃	OHC–CH ₂ –COOH	0.6 ± 0.3 (0.1–1.3)	0.6 ± 0.3 (0.1–1.0)	0.6 ± 0.3 (0.2–1.3)
4-Oxobutanoic, ω C ₄	OHC–(CH ₂) ₂ –COOH	0.9 ± 0.4 (0–1.8)	0.9 ± 0.4 (0.1–1.8)	0.8 ± 0.5 (0–1.6)
7-Oxoheptanoic, ω C ₇	OHC–(CH ₂) ₅ –COOH	0.7 ± 0.3 (0–1.2)	0.7 ± 0.3 (0–1.2)	0.5 ± 0.2 (0.3–1.0)
8-Oxoctanoic, ω C ₈	OHC–(CH ₂) ₆ –COOH	0.1 ± 0.1 (0–0.5)	0.2 ± 0.1 (0–0.5)	0.1 ± 0.1 (0–0.3)
9-Oxononanoic, ω C ₉	OHC–(CH ₂) ₇ –COOH	0.3 ± 0.1 (0.1–0.6)	0.3 ± 0.1 (0.1–0.6)	0.2 ± 0.0 (0.2–0.3)
Subtotal		8.4 ± 4.3 (1.9–19)	9.0 ± 4.4 (1.9–19)	6.6 ± 3.3 (2.7–14)
III. α-Dicarbonyls (ng m⁻³)				
Glyoxal, Gly	OHC–CHO	0.9 ± 0.7 (0.1–2.6)	1.0 ± 0.7 (0.1–2.6)	0.6 ± 0.4 (0.2–1.3)
Methylglyoxal, mGly	CH ₃ –(O)C–CHO	1.8 ± 1.5 (0–5.5)	2.1 ± 1.6 (0–5.5)	1.1 ± 1.0 (0.1–2.8)
Subtotal		2.7 ± 2.1 (0.3–7.9)	3.1 ± 2.2 (0.3–7.0)	1.7 ± 1.4 (0.3–4.1)
Total (all detected species)		242 ± 125 (75–600)	263 ± 133 (75–600)	187 ± 80 (112–349)
WSOC (μ g m ⁻³)		0.7 ± 0.3 (0.2–1.4)	0.7 ± 0.3 (0.2–1.4)	0.5 ± 0.4 (0.2–1.3)
OC (μ g m ⁻³)		1.6 ± 0.6 (0.7–2.6)	1.6 ± 0.6 (0.7–2.6)	1.3 ± 0.7 (0.7–2.5)
EC (μ g m ⁻³)		0.4 ± 0.2 (0.0–0.9)	0.4 ± 0.2 (0.1–0.9)	0.2 ± 0.2 (0.0–0.6)
PM _{2.5} (μ g m ⁻³)		22 ± 13 (3.8–62)	21 ± 12 (3.8–62)	22 ± 19 (6.2–53)
T (°C)		14 ± 2.8 (9.0–20)	15 ± 2.9 (9.0–20)	13 ± 2.3 (11–17)




Fig. 2. Molecular composition of dicarboxylic acids, ketocarboxylic acids, and α -dicarbonyls in $PM_{2.5}$ aerosols collected at Qinghai Lake, Tibet Plateau.

are comparable to those in Sonnblick ($20\text{--}76\text{ ng m}^{-3}$, Table S2), a mountain peak in the central Alps (3106 m a.s.l.) (Legrand et al., 2007), and Chichi-jima Island in North Pacific ($1.6\text{--}41\text{ ng m}^{-3}$, Table S2) (Mochida et al., 2003). Relative abundance of oxalic acid to the total diacids in the Qinghai Lake samples is $58 \pm 8\%$, followed by malonic acid (C_3 , $11 \pm 5.0\%$), succinic acid (C_4 , $6.0 \pm 2.0\%$), azelaic acid (C_9 , $5.0 \pm 2.0\%$) and phthalic acid (Ph, $3 \pm 1\%$). Such a molecular composition is similar to that in remote areas such as the western Pacific Ocean and the central Himalayan but different from those in urban regions where phthalic and/or tere-phthalic acids are more abundant than C_9 due to high emissions of anthropogenic precursors (e.g., PAHs and plasticizers) (Wang et al., 2006a; Ho et al., 2007; Cheng et al., 2013). C_9 is mainly produced from the oxidation of biogenic unsaturated fatty acids containing a double bond at the C-9 position (Kawamura and Gagosian, 1987). Relatively higher abundance of C_9 in the Qinghai Lake samples can be attributed to the abundant unsaturated fatty acids emitted from terrestrial plants and subsequent photooxidation (Kawamura and Gagosian, 1987). Averaged ratio of saturated fatty acid $C_{18:0}$ to unsaturated fatty acid $C_{18:1}$ in the $PM_{2.5}$ samples is 8.4 (calculated from the data reported by Li et al. (2013), which is much higher than that in urban area (e.g., 14 Chinese megacities 2.3 ± 0.5) (calculated from the data reported by Wang et al. (2006a)), indicating an enhanced photo-oxidation in the Tibet Plateau region.

Ketocarboxylic acids and α -dicarbonyls in the $PM_{2.5}$ samples are $8.4 \pm 4.3\text{ ng m}^{-3}$ and $2.7 \pm 2.1\text{ ng m}^{-3}$, respectively. Both classes of compounds are the precursors of dicarboxylic acids. As shown in Fig. 2, the most abundant ketocarboxylic acid is glyoxylic acid (ωC_2), comprising 55% of the total, followed by pyruvic acid (Pyr, 12%) and 3-oxobutanoic acid (ωC_4 , av. 8.0%). As shown in Table 1, glyoxal (Gly) was less abundant than methylglyoxal (mGly) ($0.9 \pm 0.7\text{ ng m}^{-3}$ versus $1.8 \pm 1.5\text{ ng m}^{-3}$), because mGly is of a stronger source (Fu et al., 2008) and OH radical oxidation with aerosol-phase mGly is slower compared to Gly (Cheng et al., 2013).

Averaged ratios of total diacid-C to OC (calculated as the ratio of the carbon concentration of all diacids to OC) in the Qinghai Lake samples are $5.1 \pm 1.7\%$, which are about three times higher than that reported for Asian megacities such as 14 Chinese cities (1.4%) (Ho et al., 2007), Tokyo, Japan (0.95%) (Kawamura and Yasui, 2005), Chennai, India (1.6%) (Pavuluri et al., 2010), and remote sites (e.g.: western Pacific, 3.2%; Jeju Island, South Korea,

3.1%) (Sempéré and Kawamura, 2003; Kundu et al., 2010b), again demonstrating more intensive photochemical oxidation nature of the Tibet Plateau atmosphere due to much stronger solar radiation.




Fig. 3. Linear fit regression for (a) temperature (T) with C_2/C_4 and C_3/C_4 ratios and (b) C_2 with SO_4^{2-} (See the abbreviations in Table 1).




Fig. 4. Comparison of mass ratios of C_6/C_9 , Ph/C_9 and F/M in the Qinghai Lake atmosphere site with those in other regions (See the abbreviations in Table 1. Data for other regions are calculated from the same documents as those in Fig. 3).

3.2. Concentrations of OC, EC and WSOC

OC and EC in the Qinghai Lake samples ranged from 0.7 to $2.6 \mu\text{g m}^{-3}$ (ave. $1.6 \pm 0.6 \mu\text{g m}^{-3}$) and 0.03 to $0.9 \mu\text{g m}^{-3}$ (ave. $0.6 \pm 0.3 \mu\text{g m}^{-3}$), respectively, which are about one order of magnitude lower than those reported for 14 Chinese cities (Cao et al., 2007). OC/EC ratio of the Qinghai Lake samples is 6.0 ± 3.9 and about 50% higher than that for Chinese urban aerosols (4.2 ± 1.0) (Cao et al., 2007). Relative abundance of WSOC to OC is $41 \pm 13\%$ for the Qinghai Lake aerosols, which is also higher than those observed in the central Himalayan (29%) (Hegde and Kawamura, 2012) and the Northwest Pacific (i.e., Gosan, Korea)

Table 2
Principal component analysis for major species in $PM_{2.5}$ aerosols from Qinghai Lake.

Compounds	Component			
	1	2	3	4
C_2	0.75	0.54	0.23	0.23
C_3	0.90	0.78	0.16	0.06
C_4	0.80	0.36	0.24	0.29
C_5	0.76	0.42	0.23	0.20
C_6	0.91	0.06	0.30	-0.16
C_9	0.90	0.07	0.18	0.15
Pyr	0.53	0.61	-0.41	0.42
ωC_2	0.67	0.43	0.23	0.24
mGly	0.54	0.67	0.20	0.27
Gly	0.80	0.41	0.15	0.33
WSOC	0.38	0.49	0.42	0.44
Levogluconan	0.12	0.12	0.17	0.88
Mg^{2+}	0.41	0.10	0.82	0.30
Ca^{2+}	0.29	0.11	0.86	0.30
K^+	0.15	0.22	0.64	0.18
SO_4^{2-}	0.17	0.68	0.28	0.12
NO_3^-	0.16	0.58	0.38	0.42
Variance (%)	61	13	8	7

Absolute values higher than 0.5 are highlighted in bold.

($30 \pm 12\%$) (Miyazaki et al., 2007), mostly due to an enhanced photochemical oxidation (Aggarwal and Kawamura, 2009).

3.3. Origins and formation mechanisms of diacids in the Qinghai Lake atmosphere

Several studies proposed that the ratios of C_2/C_4 and C_3/C_4 are indicative of aerosol aging, because they found that C_2 , C_3 and C_4 are produced by the photochemical degradation of longer-chain dicarboxylic acids, and the production of C_3 comes from the chemical oxidation of C_4 (Jung et al., 2010; Sorooshian et al., 2007b). C_2/C_4 and C_3/C_4 ratios in the Qinghai Lake samples are 11 ± 7.2 and 2.2 ± 1.3 , respectively, higher than those reported for urban aerosols such as 14 Chinese megacities (10 ± 3.1 and 0.9 ± 0.4) (Ho et al., 2007), Sapporo, Japan (3.3 and 1.3) (Aggarwal and Kawamura, 2008), marine aerosols such as Jeju Island, Korea (8.2 and 1.1) (Kundu et al., 2010b) and high alpine site in Austria (0.9 and 1.6) (Legrand et al., 2007) (Fig. S1). Pavuluri et al. (2010) found that C_2/C_4 and C_3/C_4 ratios could have strong correlation with ambient temperatures when local photooxidation is more important than long-range transport. In this study, both ratios of C_2/C_4 ($R^2 = 0.03$) and C_3/C_4 ($R^2 = 0.04$) exhibited very weak correlations with ambient temperatures (Fig. 3a), indicating that dicarboxylic acids and related SOA in the Qinghai Lake atmosphere are largely derived from long-range transport.

Table 3
Comparison of mass ratios of oxalic acid (C_2), methylglyoxal (mGly) and glyoxal to levoglucosan (Lev) in fresh $PM_{2.5}$ aerosols emitted from yak dung burning with those in ambient $PM_{2.5}$ aerosols.

Ratios	Fresh aerosols ($n = 3$)	Ambient aerosols ($n = 34$)
C_2/Lev	0.5 ± 0.3	200 ± 71
mGly/Lev	0.05 ± 0.03	2.3 ± 1.5
Gly/Lev	0.03 ± 0.01	1.1 ± 0.6

Several field observations found a robust correlation between C₂ and SO₄²⁻ and proposed a common formation pathway, i.e., in-cloud (Warneck, 2003) or aqueous-phase formation (Sullivan and Prather, 2007). In this study, we also found a strong correlation between C₂ and SO₄²⁻ (R² = 0.50) (Fig. 3b), indicating that SOA and sulfate at the continental background site share a common atmospheric process.

Maleic acid (M) can be isomerized to fumaric acid (F) under an intensive radiation (Agarwal et al., 2010). It was reported that the isomerization of M to F is suppressed under haze conditions due to weak sunlight (Hegde and Kawamura, 2012). As shown in Fig. 4, ratio of F/M (0.83 ± 0.46) in the Qinghai Lake atmosphere is higher than that reported in previous studies such as 14 Chinese cities (Ho et al., 2007) and Mt. Tai (Wang et al., 2009), probably indicating an enhanced isomerization of M to F under the Tibet Plateau strong solar radiation condition.

Adipic (C₆) and phthalic (Ph) acids are believed to be produced by the photochemical oxidation of cyclohexene and aromatic hydrocarbons such as naphthalene, which primarily originate from

anthropogenic sources (Kawamura and Ikushima, 1993). Conversely, azelaic (C₉) acid is the oxidation product of unsaturated fatty acids emitted from the biogenic sources (Kawamura and Gagosian, 1987). As a result, both ratios of C₆/C₉ and Ph/C₉ can be used to qualitatively assess the strength of anthropogenic source versus biogenic source. As seen in Fig. 4, the values of C₆/C₉ (0.1 ± 0.4) and Ph/C₉ (0.6 ± 0.2) ratios in the Qinghai Lake air are 3–10 times lower than those reported for urban, marine and mountainous atmospheres; on the contrary, C₉ to total diacids ratios of the Qinghai Lake samples are several times higher than those in the northern part of China, again demonstrating that anthropogenic sources are negligible for Qinghai Lake aerosols while biogenic sources are important.

3.4. Principal component analysis

Here we employed principal component analysis (PCA) to further explore the sources and formation mechanisms of diacids and related SOA in the Qinghai Lake atmosphere. Four components




Fig. 5. Difference in compositions of water-soluble species between the easterly samples and the westerly samples ((a) inorganic ions, (b) dicarboxylic acids, ketocarboxylic acids, and ω -dicarboxyls).

were deemed to get through the screen test, accounting for 61%, 13%, 8%, and 7% of the variance in each. As shown in Table 2, C₂–C₆, C₉ and Pyr, ωC₂, mGly and Gly showed significant correlations with the first component, while C₂, C₃, Pyr, mGly, SO₄²⁻ and NO₃⁻ well correlated with the second component. SO₄²⁻ and NO₃⁻ are regarded as the tracers of secondary aerosols and C₂ is proposed to be formed by hydroxyl radical (OH·) oxidation of various precursors including longer-chain diacids and glyoxylic acid (ωC₂) (Kawamura and Kaplan, 1987; Lim et al., 2005). Thus both components represent photochemical oxidation. However, we cannot distinguish the two components at the current stage. Component 3 presented high loadings with Mg²⁺, Ca²⁺ and K⁺ (>0.6, Table 3), representing dust source. Levoglucosan, which is a tracer of biomass burning emissions (Fu et al., 2010), had a higher loading (0.88, Table 3) with the fourth component, while all other species presented much lower loadings (<0.42, Table 3), again demonstrating the unimportance of biomass burning emissions as the source of dicarboxylic acids and related compounds at this remote site.

3.5. Backward trajectory analysis

Summertime air masses in Qinghai Lake during the sampling period were largely transported from the easterly and westerly directions (Fig. 1). Therefore, the total samples could be classified as two categories: easterly and westerly. As shown in Fig. 5a, relative abundances of SO₄²⁻ and NH₄⁺ were lower in air masses transported by the westerly than those by the easterly, but relative abundance of Ca²⁺, Mg²⁺, Na⁺ and Cl⁻ were higher in the westerly samples due to the proximity of Qaidam desert in the west. Dicarboxylic acids, ketoacids and α-dicarbonyls in the easterly air masses were more abundant than those in the westerly air masses (Table 1). Moreover, the relative abundances of smaller diacids (C₂–C₅), ωC₂ and α-dicarbonyls were higher in the easterly than in the westerly, indicating that aerosols transported from the westerly are less oxidized (Fig. 5b). The more abundant dicarboxylic acids and related compounds in the easterly samples might suggest that the easterly air masses contain more pollutants, because the eastern part to Qinghai Lake is populated area.

3.6. Comparison with fresh aerosols from yak dung burning

Since abundant C₂, mGly and Gly are found in biomass burning polumes, it has been assumed that biomass combustion can directly produce these compounds (Gao et al., 2003; Falkovich et al., 2005; Fu et al., 2008; Kundu et al., 2010a; Sorooshian et al., 2007b; Narukawa and Kawamura, 1999). Yamasoe et al. (2000) analyzed particles from direct emissions of vegetation fires and found that biomass burning could produce a certain amount of oxalic acid, accounting for 0.07 ± 0.04% and 0.04 ± 0.02% of particle mass under flaming and smoldering conditions, respectively. Yak dung is the major energy source for Tibetan in the region, thus we analyzed PM_{2.5} samples collected from the local residential area when burning yak dung for cooking in order to figure out whether biomass burning is an important source of C₂, mGly and Gly. Levoglucosan is a key tracer for biomass burning emission and chemically stable in the atmosphere (Hoffmann et al., 2010), thus the concentration ratios of C₂, mGly and Gly to levoglucosan can be used to evaluate the impact of biomass burning emission on the ambient aerosol composition. As shown in Table 3, ratio of C₂ to levoglucosan in the fresh PM_{2.5} particles is 400 times lower than that in the ambient PM_{2.5} aerosols. Likewise, the ratios of mGly and Gly to levoglucosan are 0.05 ± 0.03 and 0.03 ± 0.01 in the fresh aerosols and 30–40 times less than the ambient samples. Such significant differences between the fresh aerosols and the ambient

particles clearly demonstrate that C₂, mGly and Gly in the samples are mostly derived from photochemical oxidation with a negligible amount directly produced from biomass combustion process. Such a result is in agreement with that reported by Carlton et al. (2007), who found that volatile organic precursors emitted from biomass burning (e.g. isoprene, acetone, monoterpenes, acetylene, etc.) could contribute significantly to ambient mGly and Gly through their secondary photo-oxidation products.

4. Summary and conclusion

Dicarboxylic acids and related compounds of PM_{2.5} collected in Qinghai Lake of Tibetan Plateau during summer 2010 were characterized. C₂ is the most abundant species (58 ± 8% of total dicarboxylic acids), followed by C₃, C₄ and C₉. Concentrations of dicarboxylic acids, ketocarboxylic acids and α-dicarbonyls in the Qinghai Lake aerosols are 2–5 times lower than those in Chinese urban areas. Molecular compositions and the insignificant correlation of C₃/C₄ and C₂/C₄ ratios with temperatures demonstrate that secondary organic aerosols in the Qinghai Lake atmosphere are largely derived from long-range transport. PM_{2.5} aerosols emitted from yak dung combustion were analyzed for dicarboxylic acids and related compounds, of which results demonstrate that biomass combustion process does not directly produce significant amount of oxalic acid, Gly and mGly and these compounds in the atmosphere are mostly derived from photochemical oxidation.

Acknowledgments

This work was financially supported by the “Strategic Priority Research Program” of the Chinese Academy of Sciences (Grant No. XDA05100103, XDB05020401), and the Ministry of Science & Technology of China (2007BAC30B00, 2012BAH31B00).

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.atmosenv.2013.07.024>.

References

- Aggarwal, S.G., Kawamura, K., 2008. Molecular distributions and stable carbon isotopic compositions of dicarboxylic acids and related compounds in aerosols from Sapporo, Japan: Implications for photochemical aging during long-range atmospheric transport. *J. Geophys. Res. – Atmospheres* 113. <http://dx.doi.org/10.1029/2007JD009365>.
- Aggarwal, S.G., Kawamura, K., 2009. Carbonaceous and inorganic composition in long-range transported aerosols over northern Japan: Implication for aging of water-soluble organic fraction. *Atmos. Environ.* 43, 2532–2540.
- Agarwal, S., Aggarwal, S.G., Okuzawa, K., Kawamura, K., 2010. Size distributions of dicarboxylic acids, ketoacids, α-dicarbonyls, sugars, WSOC, OC, EC and inorganic ions in atmospheric particles over Northern Japan: implication for long-range transport of Siberian biomass burning and East Asian polluted aerosols. *Atmos. Chem. Phys.* 10, 5839–5858.
- Cao, J.J., Lee, S.C., Chow, J.C., Watson, J.G., Ho, K.F., 2007. Spatial and seasonal distributions of carbonaceous aerosols over China. *J. Geophys. Res. – Atmospheres* 112. <http://dx.doi.org/10.1029/2006JD008205>.
- Carlton, A.G., Turpin, B.J., Altieri, K.E., Seitzinger, S., Reff, A., Lim, H.-J., Ervens, B., 2007. Atmospheric oxalic acid and SOA production from glyoxal: results of aqueous photooxidation experiments. *Atmos. Environ.* 41, 7588–7602.
- Cheng, C.L., Wang, G.H., Zhou, B.H., Meng, J.J., Li, J.J., Cao, J.J., Xiao, S., 2013. Comparison of dicarboxylic acids and related compounds in aerosol samples collected in Xi'an, China during haze and clean periods. *Atmos. Environ.* (submitted for publication).
- Falkovich, A.H., Graber, E.R., Schkolnik, G., Rudich, Y., Maenhaut, W., Artaxo, P., 2005. Low molecular weight organic acids in aerosol particles from Rondonia, Brazil, during the biomass-burning, transition and wet periods. *Atmos. Chem. Phys.* 5, 781–797.
- Fu, P.Q., Kawamura, K., Pavuluri, C.M., Swaminathan, T., Chen, J., 2010. Molecular characterization of urban organic aerosol in tropical India: contributions of primary emissions and secondary photooxidation. *Atmos. Chem. Phys.* 10, 2663–2689.

- Fu, T.-M., Jacob, D.J., Wittrock, F., Burrows, J.P., Vrekoussis, M., Henze, D.K., 2008. Global budgets of atmospheric glyoxal and methylglyoxal, and implications for formation of secondary organic aerosols. *J. Geophys. Res. – Atmospheres* 113. <http://dx.doi.org/10.1029/2007JD009505>.
- Gao, S., Hegg, D.A., Hobbs, P.V., Kirchstetter, T.W., Magi, B.I., Sadilek, M., 2003. Water-soluble organic components in aerosols associated with savanna fires in southern Africa: identification, evolution, and distribution. *J. Geophys. Res. – Atmospheres* 108. <http://dx.doi.org/10.1029/2002JD002324>.
- He, N.N., Kawamura, K., 2010. Distributions and diurnal changes of low molecular weight organic acids and α -dicarbonyls in suburban aerosols collected at Mangshan, North China. *Geochem. J.* 44, e17–e22.
- Hegde, P., Kawamura, K., 2012. Seasonal variations of water-soluble organic carbon, dicarboxylic acids, ketoacids, and α -dicarbonyls in the central Himalayan aerosols. *Atmos. Chem. Phys.* 12, 6645–6665.
- Ho, K.F., Cao, J.J., Lee, S.C., Kawamura, K., Zhang, R.J., 2007. Dicarboxylic acids, ketocarboxylic acids, and dicarbonyls in the urban atmosphere of China. *J. Geophys. Res. – Atmospheres* 112. <http://dx.doi.org/10.1029/2006JD008011>.
- Ho, K.F., Lee, S.C., Ho, S.S.H., Kawamura, K., Tachibana, E., Cheng, Y., Zhu, T., 2010. Dicarboxylic acids, ketocarboxylic acids, α -dicarbonyls, fatty acids, and benzoic acid in urban aerosols collected during the 2006 Campaign of Air Quality Research in Beijing (CAREBeijing-2006). *J. Geophys. Res. – Atmospheres* 115. <http://dx.doi.org/10.1029/2009JD013304>.
- Hoffmann, D., Tilgner, A., Iinuma, Y., Herrmann, H., 2010. Atmospheric stability of levoglucosan: a detailed laboratory and modeling study. *Environ. Sci. Technol.* 44, 694–699.
- Jacobson, M.C., Hansson, H.C., Noone, K.J., Charlson, R.J., 2000. Organic atmospheric aerosols: review and state of the science. *Rev. Geophys.* 38, 267–294.
- Jung, J., Tsatsral, B., Kim, Y.J., Kawamura, K., 2010. Organic and inorganic aerosol compositions in Ulaanbaatar, Mongolia, during the cold winter of 2007 to 2008: Dicarboxylic acids, ketocarboxylic acids, and α -dicarbonyls. *J. Geophys. Res. – Atmospheres* 115. <http://dx.doi.org/10.1029/2010JD014339>.
- Kawamura, K., Gagosian, R.B., 1987. Implications of ω -oxocarboxylic acids in the remote marine atmosphere for photo-oxidation of unsaturated fatty acids. *Nature* 325, 330–332.
- Kawamura, K., Ikushima, K., 1993. Seasonal changes in the distribution of dicarboxylic acids in the urban atmosphere. *Environ. Sci. Technol.* 27, 2227–2235.
- Kawamura, K., Kaplan, I.R., 1987. Motor exhaust emissions as a primary source for dicarboxylic acids in Los Angeles ambient air. *Environ. Sci. Technol.* 21, 105–110.
- Kawamura, K., Sakaguchi, F., 1999. Molecular distributions of water soluble dicarboxylic acids in marine aerosols over the Pacific Ocean including tropics. *J. Geophys. Res. – Atmospheres* 104, 3501–3509.
- Kawamura, K., Yasui, O., 2005. Diurnal changes in the distribution of dicarboxylic acids, ketocarboxylic acids and dicarbonyls in the urban Tokyo atmosphere. *Atmos. Environ.* 39, 1945–1960.
- Kundu, S., Kawamura, K., Andreae, T.W., Hoffe, r A., Andreae, M.O., 2010a. Molecular distributions of dicarboxylic acids, ketocarboxylic acids and α -dicarbonyls in biomass burning aerosols: implications for photochemical production and degradation in smoke layers. *Atmos. Chem. Phys.* 10, 2209–2225.
- Kundu, S., Kawamura, K., Lee, M., 2010b. Seasonal variations of diacids, ketoacids, and α -dicarbonyls in aerosols at Gosan, Jeju Island, South Korea: Implications for sources, formation, and degradation during long-range transport. *J. Geophys. Res. – Atmospheres* 115. <http://dx.doi.org/10.1029/2010JD013973>.
- Latif, M.T., Peter, B., 2004. Surfactants in atmospheric aerosols. *Environ. Sci. Technol.* 38, 6501–6506.
- Legrand, M., Preunkert, S., Oliveira, T., Pio, C.A., Hammer, S., Gelencsér, A., Kasper-Giebl, A., Laj, P., 2007. Origin of C_2 – C_5 dicarboxylic acids in the European atmosphere inferred from year-round aerosol study conducted at a west–east transect. *J. Geophys. Res. – Atmospheres* 112. <http://dx.doi.org/10.1029/2006JD008019>.
- Li, J.J., Wang, G.H., Wang, X.M., Cao, J.J., Sun, T., Cheng, C.L., Meng, J.J., Hu, T.F., Liu, S.X., 2013. Abundance, composition and source of atmospheric PM_{2.5} at a remote site in Tibetan Plateau, China. *Tellus B* (submitted for publication).
- Lim, H.-J., Carlton, A.G., Turpin, B.J., 2005. Isoprene forms secondary organic aerosol through cloud processing: model simulations. *Environ. Sci. Technol.* 39, 4441–4446.
- Miyazaki, Y., Kondo, Y., Han, S., Koike, M., Kodama, D., Komazaki, Y., Tanimoto, H., Matsueda, H., 2007. Chemical characteristics of water-soluble organic carbon in the Asian outflow. *J. Geophys. Res. – Atmospheres* 112. <http://dx.doi.org/10.1029/2007JD009116>.
- Mochida, M., Kawabata, A., Kawamura, K., Hatsushika, H., Yamazaki, K., 2003. Seasonal variation and origins of dicarboxylic acids in the marine atmosphere over the western North Pacific. *J. Geophys. Res. – Atmospheres* 108. <http://dx.doi.org/10.1029/2002JD002355>.
- Narukawa, M., Kawamura, K., 1999. Distribution of dicarboxylic acids and carbon isotopic compositions in aerosols from 1997 Indonesian forest fires. *Geophys. Res. Lett.* 26, 3101–3104.
- Pavuluri, C.M., Kawamura, K., Swaminathan, T., 2010. Water-soluble organic carbon, dicarboxylic acids, ketoacids, and α -dicarbonyls in the tropical Indian aerosols. *J. Geophys. Res. – Atmospheres* 115. <http://dx.doi.org/10.1029/2009JD012661>.
- Sempéré, R., Kawamura, K., 2003. Trans-hemispheric contribution of C_2 – C_{10} , α , ω -dicarboxylic acids, and related polar compounds to water-soluble organic carbon in the western Pacific aerosols in relation to photochemical oxidation reactions. *Global Biogeochem. Cycl.* 17, 1069.
- Sorooshian, A., Lu, M.-L., Brechtel, F.J., Jonsson, H., Feingold, G., Flagan, R.C., Seinfeld, J.H., 2007a. On the source of organic acid aerosol layers above clouds. *Environ. Sci. Technol.* 41, 4647–4654.
- Sorooshian, A., Ng, N.L., Chan, A.W.H., Feingold, G., Flagan, R.C., Seinfeld, J.H., 2007b. Particulate organic acids and overall water-soluble aerosol composition measurements from the 2006 Gulf of Mexico Atmospheric Composition and Climate Study (GoMACCS). *J. Geophys. Res. – Atmospheres* 112. <http://dx.doi.org/10.1029/2007JD008537>.
- Sullivan, R.C., Prather, K.A., 2007. Investigations of the diurnal cycle and mixing state of oxalic acid in individual particles in Asian aerosol outflow. *Environ. Sci. Technol.* 41, 8062–8069.
- Wang, G.H., Kawamura, K., Cheng, C.L., Li, J.J., Cao, J.J., Zhang, R.J., Liu, S.X., Zhao, Z.Z., 2012. Molecular distribution and stable carbon isotopic composition of dicarboxylic acids, ketocarboxylic acids, and α -dicarbonyls in size-resolved atmospheric particles from Xi'an City, China. *Environ. Sci. Technol.* 46, 4783–4791.
- Wang, G.H., Niu, S.L., Liu, C., Wang, L.S., 2002. Identification of dicarboxylic acids and aldehydes of PM₁₀ and PM_{2.5} aerosols in Nanjing, China. *Atmos. Environ.* 36, 1941–1950.
- Wang, G.H., Kawamura, K., Lee, S.C., Ho, K.F., Cao, J.J., 2006a. Molecular, seasonal, and spatial distributions of organic aerosols from fourteen Chinese cities. *Environ. Sci. Technol.* 40, 4619–4625.
- Wang, G.H., Kawamura, K., Watanabe, T., Lee, S.C., Ho, K.F., Cao, J.J., 2006b. High loadings and source strengths of organic aerosols in China. *Geophys. Res. Lett.* 33. <http://dx.doi.org/10.1029/2006GL027624>.
- Wang, H.B., Kawamura, K., Yamazaki, K., 2006c. Water-Soluble dicarboxylic acids, ketoacids and dicarbonyls in the atmospheric aerosols over the southern ocean and western pacific ocean. *J. Atmos. Chem.* 53, 43–61.
- Wang, G.H., Kawamura, K., Umemoto, N., Xie, M.J., Hu, S.Y., Wang, Z.F., 2009. Water-soluble organic compounds in PM_{2.5} and size-segregated aerosols over Mount Tai in North China Plain. *J. Geophys. Res. – Atmospheres* 114. <http://dx.doi.org/10.1029/2008JD011390>.
- Warneck, P., 2003. In-cloud chemistry opens pathway to the formation of oxalic acid in the marine atmosphere. *Atmos. Environ.* 37, 2423–2427.
- Yamasoe, M.A., Artaxo, P., Miguel, A.H., Allen, A.G., 2000. Chemical composition of aerosol particles from direct emissions of vegetation fires in the Amazon Basin: water-soluble species and trace elements. *Atmos. Environ.* 34, 1641–1653.