



DR. JI CHEN (Orcid ID : 0000-0001-7026-6312)
DR. PABLO GARCIA-PALACIOS (Orcid ID : 0000-0002-6367-4761)
DR. XUHUI ZHOU (Orcid ID : 0000-0002-2038-9901)
DR. JIANWEI LI (Orcid ID : 0000-0002-0429-3627)
DR. JIANYANG XIA (Orcid ID : 0000-0001-5923-6665)
DR. SHULI NIU (Orcid ID : 0000-0002-2394-2864)
DR. KEES JAN VAN GROENIGEN (Orcid ID : 0000-0002-9165-3925)

Article type : Primary Research Articles

Differential responses of carbon-degrading enzymes activities to warming:
implications for soil respiration

Running title: Carbon-degrading enzyme responses to warming

Ji Chen^{1,2,3*} | Yiqi Luo^{4,5} | Pablo García-Palacios⁶ | Junji Cao^{2,7*} | Marina Dacal⁶ | Xuhui Zhou^{8,9} | Jianwei Li^{10*} | Jianyang Xia⁸ | Shuli Niu¹¹ | Huiyi Yang¹² | Shelby Shelton¹³ | Wei Guo¹⁴ | Kees Jan van Groenigen¹⁵

¹Center for Ecological and Environmental Sciences, Key Laboratory for Space Bioscience & Biotechnology, Northwestern Polytechnical University, Xi'an, 710072, China;

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/gcb.14394

This article is protected by copyright. All rights reserved.

Accepted Article

²State Key Laboratory of Loess and Quaternary Geology (SKLLQG), and Key Laboratory of Aerosol Chemistry and Physics, Institute of Earth Environment, Chinese Academy of Sciences, Xi'an, 710061, China;

³Aarhus University Centre for Circular Bioeconomy, Department of Agroecology, Aarhus University, Blichers Allé 20, 8830 Tjele, Denmark.

⁴Center for Ecosystem Science and Society, Department of Biological Sciences, Northern Arizona University, Arizona, 86011 USA;

⁵Department for Earth System Science, Tsinghua University, Beijing 100084, China.

⁶Departamento de Biología y Geología, Física y Química Inorgánica y Analítica, Área de Biodiversidad y Conservación, Universidad Rey Juan Carlos, 28933 Móstoles, Spain;

⁷Institute of Global Environmental Change, Xi'an Jiaotong University, Xi'an, 710049, China;

⁸Center for Global Change and Ecological Forecasting, Tiantong National Field Observation Station for Forest Ecosystem, School of Ecological and Environmental Sciences, East China Normal University, Shanghai, 200062, China.

⁹Shanghai Institute of Pollution Control and Ecological Security, Shanghai 200437, China.

¹⁰Department of Agriculture and Environmental Sciences, Tennessee State University, Nashville, 37209, USA;

¹¹Synthesis Research Center of Chinese Ecosystem Research Network, Key Laboratory of Ecosystem Network Observation and Modeling, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing, 100101, China;

¹²College of engineering mathematics and physical sciences, University of Exeter, EX4 4QF, UK

¹³Department of Emergency Medicine, University of Colorado Denver, Denver, CO, 80045 USA;

¹⁴Department of Earth and Environmental Sciences, Xi'an Jiaotong University, Xi'an 710049, China;

¹⁵Geography, College of Life and Environmental Sciences, University of Exeter, Exeter, EX4 4QG, UK.

*Corresponding author:

Ji Chen, chenji@ieecas.cn, Tel: +86-29-62336233, Fax: +86-29-62336234.

Junji Cao, cao@loess.llqg.ac.cn, Tel: +86-29-62336233, Fax: +86-29-62336234.

Jianwei Li, jli2@tnstate.edu, Tel: +1-615-963-1523, Fax: +1-615-963-1523.

This article is protected by copyright. All rights reserved.

Abstract

Extracellular enzymes catalyze rate-limiting steps in soil organic matter decomposition, and their activities (EEAs) play a key role in determining soil respiration (SR). Both EEAs and SR are highly sensitive to temperature, but their responses to climate warming remain poorly understood. Here, we present a meta-analysis on the response of soil cellulase and ligninase activities and SR to warming, synthesizing data from 56 studies. We found that warming significantly enhanced ligninase activity by 21.4% but had no effect on cellulase activity. Increases in ligninase activity were positively correlated with changes in SR, while no such relationship was found for cellulase. The warming response of ligninase activity was more closely related to the responses of SR than a wide range of environmental and experimental methodological factors. Furthermore, warming effects on ligninase activity increased with experiment duration. These results suggest that soil microorganisms sustain long term increases in SR with warming by gradually increasing the degradation of the recalcitrant carbon pool.

KEYWORDS

Extracellular enzyme activity, Decomposition, Soil microorganisms, Soil respiration, Recalcitrant carbon pool, Ligninase activity, Cellulase activity, Global warming

1 | INTRODUCTION

The average global surface temperature is predicted to increase between 1 and 4°C by the end of the 21st century (Collins & Knutti, 2013, O'Neill et al., 2017). Rising temperatures have cascading impacts on ecosystem carbon (C) budgets, and these can cause both positive and negative C cycle–climate feedbacks (Carey et al., 2016, Chen et al., 2016a, Chen et al., 2017a, Karhu et al., 2014, Paustian et al., 2016, Peñuelas et al., 2017, Yang et al., 2018). Soil respiration (SR) represents the largest C flux from soils to the atmosphere (Bradford et al., 2016, Tucker, Bell, Pendall, & Ogle, 2013), and is primarily driven by the microbial decomposition of soil organic matter (SOM).

However, we know little about the mechanisms underlying the response of SR to climate warming (Chen et al., 2016b, Conant et al., 2011, van Gestel et al., 2018).

Specifically, there is a lack of information regarding the degree to which soil extracellular enzymes (EEs), which catalyze the rate-limiting step in SOM decomposition (Allison, Wallenstein, & Bradford, 2010a, Jing et al., 2014, Sinsabaugh, 2010, Stone et al., 2012), are affected by warming. These enzymes, primarily produced by microbes, are considered proximate agents of SR because they lower the activation energy of key reactions and speed up the breakdown of polymers (Chen et al., 2017b, Chen et al., 2018, Janssens et al., 2010, Suseela, Tharayil, Xing, & Dukes, 2014).

Although the rates at which these enzymes are produced and degraded are sensitive to temperature (Allison & Treseder, 2008, German, Marcelo, Stone, & Allison, 2012, Papanikolaou, Britton, Helliwell, & Johnson, 2010, Steinweg, Dukes, Paul, & Wallenstein, 2013), it is still unclear how warming responses of enzymes affect SR.

This article is protected by copyright. All rights reserved.

Cellulose and lignin are the two most abundant SOM compounds, and microbially mediated decomposition of these materials composes a main source of SR (Carreiro, Sinsabaugh, Repert, & Parkhurst, 2000, Chen et al., 2018, Janssens et al., 2010, Waldrop et al., 2004). Cellulose and hemicellulose comprise the main composition of primary plant cell walls. Hydrolysis of cellulose and hemicellulose is mainly catalyzed by cellulase, including β -1,4-glucosidase (BG), β -1,4-xylosidase (BX) and β -D-cellobiosidase (CBH) (Carreiro, Sinsabaugh, Repert, & Parkhurst, 2000, Chen et al., 2017b, Jian et al., 2016). The aromatic C polymer lignin is found in secondary plant cell walls, where it covers and shields cellulose from microbial decay. Oxidation and degradation of phenolic-containing recalcitrant compounds are facilitated by ligninase, that is, peroxidase (PER), phenol oxidase (PO) and polyphenol oxidase (PPO) (Dashtban, Schraft, Syed, & Qin, 2010, Romero-Olivares, Allison, & Treseder, 2017, Sinsabaugh et al., 2008, Zhou et al., 2012). The critical roles of cellulase and ligninase in mediating SOM decomposition suggest that climate warming may affect SR through its effects on EEAs, yet we still lack direct evidence.

Cellulase and ligninase are synthesized by specific groups of microorganisms (Burns et al., 2013, Carreiro, Sinsabaugh, Repert, & Parkhurst, 2000, Wang et al., 2012), and it may take years for microbial communities to adapt to environmental changes (DeAngelis et al., 2015). Thus, responses of cellulase and ligninase activities to warming may vary over time. Because warming methods differ in their effects on soil temperature and moisture (Chen et al., 2015, Lu et al., 2013), soil microbial community (Chen et al., 2015) and belowground C allocation (Rustad et al., 2001, This article is protected by copyright. All rights reserved.

Accepted Article

Schindlbacher et al., 2015), they may differ in their effects on EEAs as well. Including cellulase and ligninase activities in soil C models may improve future predictions of soil C stocks (Ali et al., 2015, Luo, Chen, Chen, & Feng, 2017, Moorhead, Sinsabaugh, Hill, & Weintraub, 2016). However, warming effects on cellulase and ligninase activities as well as the underlying mechanisms are still unclear.

To address this knowledge gap, we conducted a meta-analysis of the responses of cellulase and ligninase activities to warming and their links with SR responses. More specifically, our study seeks : (1) to quantify the effects of warming on cellulase and ligninase activities, (2) to investigate the factors affecting the responses of cellulase and ligninase activities to warming, and (3) to test whether the responses of cellulase and ligninase activities to warming are linked with changes in SR.

2 | MATERIALS AND METHODS

2.1 | Data collection

We extracted results for enzyme activities of ligninase and cellulase under warming experiments conducted in the field. We used Web of Science (<http://apps.webofknowledge.com/>), Google Scholar (<http://scholar.google.com/>) and China National Knowledge Infrastructure (<http://www.cnki.net/>) for an exhaustive search of journal articles published before June 2018, using the following key words : (1) “climate change” or “experimental warming” or “elevated temperature” and (2) “cellulase”, or “ligninase”, or “glucosidase”, or “xylosidase”, or “cellobiosidase”, or

This article is protected by copyright. All rights reserved.

“peroxidase”, or “phenol oxidase”, or “polyphenol oxidase”, and (3) “terrestrial” or “soil” or “land”.

To be included in our dataset, experiments had to meet several criteria: (1) the warming treatment lasted at least one year; (2) vegetation, soil physicochemical parameters and climate were similar between control and warming treatments; (3) sample size and standard deviations were reported; (4) warming protocols (i.e., warming method, warming magnitude, warming time and warming season) were clearly described. All studies in our dataset measured enzyme activity for warmed and control soils at the same incubation temperature (i.e., temperature differences between treatments occurred only in the field, and not during the incubation). As such, differences in enzyme activity between warmed and control soils were not related to the temperature sensitivity of enzymes, but reflect warming effects on enzyme production by soil microbes. We found 56 articles that met our requirements (see Dataset and Figure S1).

For each study in our dataset, we extracted information on cellulase and ligninase activities (Table S1). If a paper reported multiple warming responses (e.g. in multi-factor experiments, or studies applying more than one warming protocol), each experiment was included separately in our dataset. If one paper reported two or three kinds of cellulase or ligninase, then their sum values were considered as the overall responses of cellulase and ligninase activities. We also recorded a wide range of environmental variables, including latitude, longitude, elevation, climatic variables (mean annual temperature (MAT), mean annual precipitation (MAP)), sampling date, This article is protected by copyright. All rights reserved.

Accepted Article

sampling temperature, vegetation type (<http://www.worldclim.org/>) and soil type (<http://www.fao.org/about/en/>). Regarding the warming protocols, we recorded the magnitude (i.e., the average temperature difference between the warming and control plots), duration (in years) and methods (open top chamber (OTC), infrared heater (IH), green house (GH), heating cable, and curtain). We also recorded SR, soil C:N, microbial biomass, and the ratio of fungal to bacterial abundance for both control and warming treatments when these data were reported. When warming responses of SR were not available, we used responses of heterotrophic respiration or weight loss in litter bag experiments as proxy values. To extract data from figures we used Engauge Digitizer 4.1 (<http://digitizer.sourceforge.net>). When some critical information was not reported in the article, we tried to obtain this information by contacting the corresponding author.

2.2 | Data analysis

We used meta-analysis to evaluate the effects of warming on cellulase, ligninase, individual enzyme activity and other ancillary variables (García-Palacios et al., 2014, Hedges, Gurevitch, & Curtis, 1999, Van Groenigen et al., 2014, Zhao et al., 2017). The effects of warming on EEAs were evaluated using the natural log of the response ratio ($\ln R$):

$$\ln R = \ln \left(\frac{\overline{X}_w}{\overline{X}_c} \right) = \ln(\overline{X}_w) - \ln(\overline{X}_c) \quad (1),$$

with \overline{X}_w and \overline{X}_c as the arithmetic mean concentrations in the warming and control treatments, respectively. The variances (v) of $\ln R$ were calculated by:

$$v = \frac{s_w^2}{n_w \overline{X}_w^2} + \frac{s_c^2}{n_c \overline{X}_c^2} \quad (2),$$

with n_w and n_c as the number of replicates, and S_w and S_c as the *SDs* for warming and control treatments, respectively.

The overall effect and the 95% confidence interval were calculated using the “rma.mv” function in the R-package “metafor” (Viechtbauer, 2010). Because incubation temperature for enzyme measurements varied among studies, we included “Incubation temperature” as a random factor in the meta-analysis. Because several papers contributed more than one response ratio, we also included the variable “paper” as a random factor (Chen et al., 2018, Terrer et al., 2016, van Groenigen et al., 2017). The effects of warming were considered significant if the 95% confidence interval did not overlap with zero. The results for the analyses on $\ln R$ were back-transformed and reported as percentage change with warming (that is, $100 \times (e^{\ln R} - 1)$) to ease interpretation.

The meta-analytic models were selected by using the same approach as in Chen et al. (2018), Terrer et al. (2016) and van Groenigen et al. (2017). Briefly, we analyzed all potential combinations of the studied factors in a mixed-effects meta-regression model using the “glmulti” package in R (Bangert-Drowns, Hurley, & Wilkinson, 2004, Calcagno & de Mazancourt, 2010). The importance of a specific predictor was expressed as the sum of Akaike weights for models that included this factor, which can be considered as the overall support for each variable across all models. A cut-off of 0.8 was set to differentiate between important and non-essential predictors.

3 | RESULTS

Across the whole dataset, warming significantly enhanced ligninase activity by an average of 21.4%. Specifically, warming significantly increased activities of PER by 18.4%, PO by 13.5% and PPO by 28.6%. In contrast, warming had no effect on cellulase activity (Figure 1a), or any of the individual cellulase enzymes BG, BX and CBH. The responses of cellulase and ligninase activities to warming were normally distributed (Figure 1b and 1c), and they were independent of the sample size (Figure S2).

None of variables tested for the effects of warming on cellulase activity reached the threshold value (0.8) of the summed Akaike weights (Figure 2a). In contrast, effects of warming on ligninase activity were best explained by warming duration and warming method (Figure 2b). Linear regression analysis confirmed that $\ln R$ of ligninase activity

was positively correlated with warming duration, while no such relationship was found for cellulase activity (Figure 3a and 3b). Regarding warming methods, warming did not affect cellulase activity for any of the warming methods (Figure 3a). In contrast, OTC, GH and IH significantly increased ligninase activity by 15.5%, 31.4% and 22.3%, respectively; while cables had no effect on ligninase activity (Figure 3b).

Warming significantly increased microbial biomass specific ligninase activity (i.e. the ratio of ligninase activity to total microbial abundance) by 40.6% (Figure S3a). This increase was weakly positively correlated with warming-induced changes in the ratio of fungal to bacterial abundance (Figure S3b). Finally, our analyses suggest that warming had stronger positive effects on biomass specific ligninase activity for long-term than short-term studies, while this relationship was not observed for biomass specific cellulase activity (Figure S4a and S4b).

Warming on average increased SR by 15.8% (95% CI: 6.3% - 26.1%) in our dataset. We found no relationship between the responses of cellulase activity and the responses of SR to warming (Figures 4a). However, the warming response of SR was positively correlated with the response of ligninase activity and the positive relationship held when analyzed for PER, PO and PPO individually (Figures 4b and S5). To compare the relative importance of cellulase and ligninase activities in explaining the response of SR to warming, we limited our model selection analysis to studies that simultaneously reported the effects of warming on cellulase and ligninase activities and SR. Effects of warming on SR were best predicted by the responses of ligninase activity over a wide range of ecosystem types, climatic variables and warming

This article is protected by copyright. All rights reserved.

Accepted Article

protocols (Figure 4c). Experiment duration had no significant impact on SR responses to warming, either in the subset of studies that reported responses of both enzymes (Figure 4c), or across the entire dataset.

4 | DISCUSSION

Our results show that warming significantly enhanced ligninase activity, and that warming responses are positively correlated with warming duration. In contrast, warming does not affect cellulase activity. Why does warming have differential effects on cellulase and ligninase activities? We propose three possible mechanisms. First, the enzyme responses reflect warming-induced changes in substrate availability. Enzyme activity can be described by the Michaelis–Menten relationship, which primarily depends on substrate availability (Davidson & Janssens, 2006, Sinsabaugh et al., 2008). Initial stimulation of SR by warming depletes easily hydrolysable substrates (Allison, McGuire, & Treseder, 2010b, Luo, Wan, Hui, & Wallace, 2001), limiting the positive response of cellulase activity to increasing temperatures (Davidson & Janssens, 2006, Stone et al., 2012, Weedon, Aerts, Kowalchuk, & van Bodegom, 2014). At the same time, warming-induced declines in easily hydrolysable C pools can lead to microbial C starvation (Crowther & Bradford, 2013, Fenner et al., 2006, Melillo et al., 2017, Metcalfe, 2017). Under these circumstances, soil microbial communities may adapt to utilize previously inaccessible recalcitrant C pools to fuel their metabolic activities. Microbial utilization of recalcitrant substrates such as phenol requires depolymerization, a process catalyzed by ligninase (De Gonzalo, Colpa, Habib, & Fraaije, 2016, Jassey et al., 2012, Sinsabaugh, 2010).

This article is protected by copyright. All rights reserved.

Accepted Article

Second, warming may increase ligninase activity through its effect on soil N availability. Warming-induced redistribution of N from soils to vegetation could progressively lead to microbial N limitation, particularly in high C:N regions (Bai et al., 2013, Beier et al., 2008, Melillo et al., 2011). In that case, soil microorganisms are expected to invest C and energy to acquire N through decomposition of N-containing molecules (Chen et al., 2017b, Sinsabaugh et al., 2008), which are often physically or chemically protected by other aromatic macromolecules such as lignin (Hobbie, 2008, Weedon et al., 2012, Zhao et al., 2014). This explanation is supported by the positive correlation between warming effects on ligninase activity and soil C:N, while no clear relationship is found for the responses of cellulase activity (Figure S6). Finally, warming-induced changes in soil microclimate (Domínguez et al., 2017, Zhou et al., 2013), fresh C input (Bhattacharyya et al., 2013a, Xue et al., 2016, Yin et al., 2013) and plant community composition (Kardol, Cregger, Campany, & Classen, 2010, Steinauer et al., 2015) can all cause substantial changes in microbial communities as well.

Increased ligninase production with warming might reflect shifts in the microbial community composition. Indeed, several studies suggest that warming-induced changes in soil microbial community composition cause differential responses of cellulase and ligninase activities (DeAngelis et al., 2015, Pold, Grandy, Melillo, & DeAngelis, 2017). This explanation is also consistent with studies showing that fungi are main contributors to ligninase production (De Gonzalo, Colpa, Habib, & Fraaije, 2016, Kinnunen, Maijala, JArvinen, & Hatakka, 2017), and that experimental warming increases fungal abundance (A'Bear, Jones, Kandeler, & Boddy, 2014, Delarue et al.,

2015). However, warming may also directly or indirectly cause physiological adaptation of soil microorganisms to increase enzyme production (Manzoni et al., 2012, Nie et al., 2013, Schindlbacher et al., 2015), even when warming decreases total microbial biomass (Pold, Grandy, Melillo, & DeAngelis, 2017, Sistla & Schimel, 2013, Sorensen et al., 2018). This is consistent with recent findings that experimental warming tends to decrease microbial C use efficiency (Manzoni et al., 2012, Tucker, Bell, Pendall, & Ogle, 2013).

Why does the effect of warming on ligninase activity increase over time? Soil microorganisms can adjust their community composition or alter their C utilization strategies to adapt to warming, but it requires several years or even decades for significant changes in their community composition to occur (DeAngelis et al., 2015, Feng et al., 2017, Rousk, Smith, & Jones, 2013). Furthermore, warming-induced N limitation may take several years to manifest (Bai et al., 2013, Melillo et al., 2011). In addition, long-term warming could also restructure plant community and alter litter quality towards decay resistance (e.g. high lignin content) (Melillo et al., 2011, Talbot, Yelle, Nowick, & Treseder, 2012), thereby promoting the microbial production of ligninase.

Regardless of the mechanism underlying the differential warming response of ligninase and cellulase, our results suggest that warming-induced shifts in cellulase and ligninase activities could help to sustain long-term increases in SR with warming (Lin, Zhu, & Cheng, 2015, Romero-Olivares, Allison, & Treseder, 2017, Souza et al., 2017). This is because warming responses of ligninase activity exert far larger control over SR

This article is protected by copyright. All rights reserved.

than a broad range of environmental and experimental variables. These results suggest that responses of SR to warming are largely modulated by a single group of lignin-modifying enzymes, which contributes to sustained positive responses of SR to long-term climate warming.

Warming methods constituted the second important predictor of the warming effects on ligninase activity. Cables only warm soils and are reported to have negative effects on microbial biomass, litter inputs and root exudates (Rustad et al., 2001, Schindlbacher et al., 2015). Similarly, a recent meta-analysis shows that cables generally decrease total microbial, fungal and bacterial abundance, while other warming methods increase microbial abundance (Chen et al., 2015). We hypothesize that these negative responses suppressed microbial activity and microbial enzymatic production (Chen et al., 2015, Hanson et al., 2017). In addition, high warming magnitude and large reductions in soil moisture in cable experiments may decrease microbial C use efficiency (Schindlbacher et al., 2011, Schindlbacher et al., 2012), which could potentially suppress microbial cellulase and ligninase production.

Model projections of soil C dynamics often lack representation of EEAs regulated SOM decomposition (Davidson & Janssens, 2006, Luo et al., 2016, Wieder, Bonan, & Allison, 2013). However, our finding that warming-induced shifts in cellulase and ligninase activities may facilitate sustained increases in SR under long-term climate warming, highlights the need for a closer integration of enzymatic decomposition into soil biogeochemical models. Unfortunately, responses of SR and EEAs to long-term climate warming remain understudied, since experiment duration is often constrained

This article is protected by copyright. All rights reserved.

by funding availability. If the relationship between ligninase and warming duration holds across a wide range of land ecosystems, our results suggest that ecosystem climate-carbon feedbacks could be stronger than previously assumed.

ACKNOWLEDGEMENTS

Firstly, we would like to thank the authors whose work is included in this meta-analysis, especially those who supplied us with additional data. Secondly, we would like to thank Robert Sinsabaugh and three anonymous reviewers for their valuable comments on an earlier version of this manuscript. This study was supported by the National Natural Science Foundation of China (41701292), China Postdoctoral Science Foundation (2017M610647, 2018T111091), the Natural Science Basic Research Plan in Shaanxi Province (2017JQ3041), the State Key Laboratory of Loess and Quaternary Geology (SKLLQG1602), the Key Laboratory of Aerosol Chemistry and Physics (KLACP-17-02), Institute of Earth Environment, Chinese Academy of Sciences. Contributions from Dr. Luo's Eco-lab to this study were financially supported by United States Department of Energy grant DE-SC00114085, and US National Science Foundation grants EF 1137293 and OIA-1301789. This work was also supported by NSFC-Yunnan United fund (U1302267) and the National Science Fund for Distinguished Young Scholars (31325005).

DATA ACCESSIBILITY

The data associated with this paper is available from the online supplementary file.

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

CONFLICT of INTEREST

The authors declare no conflict of interest.

This article is protected by copyright. All rights reserved.

ORCID

Ji Chen <http://orcid.org/0000-0001-7026-6312>

Pablo Garcia-Palacios <http://orcid.org/0000-0002-6367-4761>

Xuhui Zhou <http://orcid.org/0000-0002-2038-9901>

Jianwei Li <http://orcid.org/0000-0002-0429-3627>

Jianyang Xia <http://orcid.org/0000-0001-5923-6665>

Shuli Niu <http://orcid.org/0000-0002-2394-2864>

Kees Jan van Groenigen <http://orcid.org/0000-0002-9165-3925>

References

Papers indicated with * are included in our meta-analysis.

*A'bear, A. D., Jones, T. H., Kandeler, E., & Boddy, L. (2014). Interactive effects of temperature and soil moisture on fungal-mediated wood decomposition and extracellular enzyme activity. *Soil Biology and Biochemistry*, 70, 151-158. doi:10.1016/j.soilbio.2013.12.017.

Ali, R. S., Ingwersen, J., Demyan, M. S., Funkuin, Y. N., Wizemann, H. D., Kandeler, E., & Poll, C. (2015). Modelling in situ activities of enzymes as a tool to explain seasonal variation of soil respiration from agro-ecosystems. *Soil Biology and Biochemistry*, 81, 291-303. doi:10.1016/j.soilbio.2014.12.001.

Allison, S. D., Wallenstein, M. D., & Bradford, M. A. (2010a). Soil-carbon response to warming dependent on microbial physiology. *Nature Geoscience*, 3, 336-340. doi:10.1038/Ngeo846.

*Allison, S. D., Mcguire, K. L., & Treseder, K. K. (2010b). Resistance of microbial and soil properties to warming treatment seven years after boreal fire. *Soil Biology and Biochemistry*, 42, 1872-1878. doi:10.1016/j.soilbio.2010.07.011.

Allison, S. D., & Treseder, K. K. (2008). Warming and drying suppress microbial activity and carbon cycling in boreal forest soils. *Global Change Biology*, 14, 2898-2909. doi:10.1111/j.1365-2486.2008.01716.x.

This article is protected by copyright. All rights reserved.

- *Bai C. (2011). The Effects of Experimental warming and nitrogen addition on soil properties. *Inner Mongolia Agricultural University*.
http://www.wanfangdata.com.cn/details/detail.do?_type=degree&id=D341262. (In Chinese with English abstract)
- Bai, E., Li, S., Xu, W., Li, W., Dai, W., & Jiang, P. (2013). A meta-analysis of experimental warming effects on terrestrial nitrogen pools and dynamics. *New Phytologist*, 199, 441-451.
doi:10.1111/nph.12252.
- Bangert-Drowns, R. L., Hurley, M. M., & Wilkinson, B. (2004). The effects of school-based writing-to-learn interventions on academic achievement: A meta-analysis. *Review of Educational Research*, 74, 29-58. doi:10.3102/00346543074001029.
- Beier, C., Emmett, B. A., Peñuelas, J., Schmidt, I. K., Tietema, A., Estiarte, M., . . . Gorissen, A. (2008). Carbon and nitrogen cycles in European ecosystems respond differently to global warming. *Science of the Total Environment*, 407, 692-697. doi:10.1016/j.scitotenv.2008.10.001.
- *Bhattacharyya, P., Roy, K. S., Neogi, S., Dash, P. K., Nayak, A. K., Mohanty, S., . . . Rao, K. S. (2013a). Impact of elevated CO₂ and temperature on soil C and N dynamics in relation to CH₄ and N₂O emissions from tropical flooded rice (*Oryza sativa* L.). *Science of the Total Environment*, 461, 601-611. doi:10.1016/j.scitotenv.2013.05.035.
- *Bhattacharyya, P., Roy, K. S., Neogi, S., Manna, M. C., Adhya, T. K., Rao, K. S., & Nayak, A. K. (2013b). Influence of elevated carbon dioxide and temperature on belowground carbon allocation and enzyme activities in tropical flooded soil planted with rice. *Environmental Monitoring and Assessment*, 185, 8659-8671. doi:10.1007/s10661-013-3202-7.
- Bradford, M. A., Wieder, W. R., Bonan, G. B., Fierer, N., Raymond, P. A., & Crowther, T. W. (2016). Managing uncertainty in soil carbon feedbacks to climate change. *Nature Climate Change*, 6, 751-758. doi:10.1038/nclimate3071.
- Burns, R. G., Deforest, J. L., Marxsen, J., Sinsabaugh, R. L., Stromberger, M. E., Wallenstein, M. D., . . . Zoppini, A. (2013). Soil enzymes in a changing environment: Current knowledge and future directions. *Soil Biology and Biochemistry*, 58, 216-234. doi:10.1016/j.soilbio.2012.11.009.
- Calcagno, V., & De Mazancourt, C. (2010). glmulti: an R package for easy automated model selection with (generalized) linear models. *Journal of Statistical Software*, 34, 1-29.
<http://www.jstatsoft.org/article/view/v034i12>.
- Carey, J. C., Tang, J., Templer, P. H., Kroeger, K. D., Crowther, T. W., Burton, A. J., . . . Heskell, M. A. (2016). Temperature response of soil respiration largely unaltered with experimental warming. *Proceedings of the National Academy of Sciences*, 113, 13797-13802.
doi:10.1073/pnas.1605365113.

- Carreiro, M., Sinsabaugh, R., Repert, D., & Parkhurst, D. (2000). Microbial enzyme shifts explain litter decay responses to simulated nitrogen deposition. *Ecology*, 81, 2359-2365. doi:10.1890/0012-9658(2000)081[2359:MESELD]2.0.CO;2.
- Chen, J., Luo, Y., Xia, J., Jiang, L., Zhou, X., Lu, M., . . . Cao, J. (2015). Stronger warming effects on microbial abundances in colder regions. *Scientific Reports*, 5, 18032. doi:10.1038/srep18032.
- Chen, J., Luo, Y., Xia, J., Shi, Z., Jiang, L., Niu, S., . . . Cao, J. (2016a). Differential responses of ecosystem respiration components to experimental warming in a meadow grassland on the Tibetan Plateau. *Agricultural and Forest Meteorology*, 220, 21-29. doi:10.1016/j.agrformet.2016.01.010.
- Chen, J., Luo, Y., Xia, J., Wilcox, K. R., Cao, J., Zhou, X., . . . Wang, R. W. (2016b). Warming effects on ecosystem carbon fluxes are modulated by plant functional types. *Ecosystems*, 1-12. doi:10.1007/s10021-016-0035-6.
- Chen, J., Zhou, X., Hruska, T., Cao, J., Zhang, B., Liu, C., . . . Wang, P. (2017a) Asymmetric diurnal and monthly responses of ecosystem carbon fluxes to experimental warming. *CLEAN – Soil, Air, Water*, 45, 1600557. doi:10.1002/clen.201600557.
- Chen, J., Luo, Y., Li, J., Zhou, X., Cao, J., Wang, R. W., . . . Walker, L. M. (2017b). Costimulation of soil glycosidase activity and soil respiration by nitrogen addition. *Global Change Biology*, 23, 1328-1337. doi:10.1111/gcb.13402.
- Chen, J., Luo, Y., Van Groenigen, K. J., Hungate, B. A., Cao, J., Zhou, X., & Wang, R. W. (2018) A keystone microbial enzyme for nitrogen control of soil carbon storage. *Science Advances*. doi:10.1126/sciadv.aag1689.
- *Chen, S. T., Sang, L., Zhang, X., & Hu, Z. H. (2016). Effects of warming and straw application on soil respiration and enzyme activity in a winter wheat cropland. *Environmental Science*, 37, 703-709. doi:10.13227/j.hjx.2016.02.040. (In Chinese with English abstract)
- *Chen, X. L., Wang, G. X., & Yang, Y. (2015). Response of soil surface enzyme activities to short-term warming and litter decomposition in a Mountain forest. *Acta Ecologica Sinica*. doi:10.5846/stxb201312182982. (In Chinese with English abstract)
- Collins, M., & Knutti, R. (2013). Long-term climate change: projections, commitments and irreversibility. In *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (Stocker, T.F., et al., eds)*, p. 1054. Cambridge, UK and New York, NY, USA: Cambridge University Press. <http://pure.iiasa.ac.at/id/eprint/10551/>.

- Accepted Article
- Conant, R. T., Ryan, M. G., Agren, G. I., Birge, H. E., Davidson, E. A., Eliasson, P. E., . . . Hopkins, F. M. (2011). Temperature and soil organic matter decomposition rates—synthesis of current knowledge and a way forward. *Global Change Biology*, 17, 3392-3404. doi:10.1111/j.1365-2486.2011.02496.x.
- Crowther, T. W., & Bradford, M. A. (2013). Thermal acclimation in widespread heterotrophic soil microbes. *Ecology Letters*, 16, 469-477. doi:10.1111/ele.12069.
- Dashtban, M., Schraft, H., Syed, T. A., & Qin, W. (2010). Fungal biodegradation and enzymatic modification of lignin. *International Journal of Biochemistry and Molecular Biology*, 1, 36. doi:PMC3180040.
- Davidson, E. A., & Janssens, I. A. (2006). Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature*, 440, 165-173. doi:10.1038/nature04514.
- De Gonzalo, G., Colpa, D. I., Habib, M. H., & Fraaije, M. W. (2016). Bacterial enzymes involved in lignin degradation. *Journal of Biotechnology*, 236, 110-119. doi:10.1016/j.jbiotec.2016.08.011.
- Deangelis, K. M., Pold, G., Topçuoğlu, B. D., Van Diepen, L. T., Varney, R. M., Blanchard, J. L., . . . Frey, S. D. (2015). Long-term forest soil warming alters microbial communities in temperate forest soils. *Frontiers in Microbiology*, 6, 104. doi:10.3389/fmicb.2015.00104.
- *Delarue, F., Gogo, S., Buttler, A., Bragazza, L., Jasse, V. E. J., Bernard, G., & Laggoun-Defarge, F. (2014). Indirect effects of experimental warming on dissolved organic carbon content in subsurface peat. *Journal of Soils and Sediments*, 14, 1800-1805. doi:10.1007/s11368-014-0945-x.
- *Delarue, F., Buttler, A., Bragazza, L., Grasset, L., Jasse, V. E. J., Gogo, S., & Laggoun-Defarge, F. (2015). Experimental warming differentially affects microbial structure and activity in two contrasted moisture sites in a Sphagnum-dominated peatland. *Science of the Total Environment*, 511, 576-583. doi:10.1016/j.scitotenv.2014.12.095.
- *Domínguez, M. T., Holthof, E., Smith, A. R., Koller, E., & Emmett, B. A. (2017). Contrasting response of summer soil respiration and enzyme activities to long-term warming and drought in a wet shrubland (NE Wales, UK). *Applied Soil Ecology*, 110, 151-155. doi:10.1016/j.apsoil.2016.11.003.
- Feng, W., Liang, J., Hale, L. E., Jung, C. G., Chen, J., Zhou, J., . . . Luo, Y. (2017). Enhanced decomposition of stable soil organic carbon and microbial catabolic potentials by long-term field warming. *Global Change Biology*, 23, 4765-4776. doi:10.1111/gcb.13755.

- *Fenner, N., Freeman, C., Lock, M. A., Harmens, H., Reynolds, B., & Sparks, T. (2006). Interactions between elevated CO₂ and warming could amplify DOC exports from peatland catchments. *Environmental Science & Technology*, 41, 3146-3152. doi:10.1021/Es061765v.
- *Gao, J., Wang, E., Ren, W., Liu, X., Chen, Y., Shi, Y., & Yang, Y. (2017). Effects of simulated climate change on soil microbial biomass and enzyme activities in young Chinese fir (*Cunninghamia lanceolata*) in subtropical China. *Acta Ecologica Sinica*, 37, 272-278. doi:10.1016/j.chnaes.2017.02.007. (In Chinese with English abstract)
- García-Palacios, P., Vandegehuchte, M. L., Ashley, Shaw E., Dam, M., Post, K. H., Ramirez, K. S., . . . Wall, D. H. (2014). Are there links between responses of soil microbes and ecosystem functioning to elevated CO₂, N deposition and warming? A global perspective. *Global Change Biology*, 21, 1590-1600. doi:10.1111/gcb.12788.
- German, D. P., Marcelo, K. R. B., Stone, M. M., & Allison, S. D. (2012). The Michaelis-Menten kinetics of soil extracellular enzymes in response to temperature: a cross-latitudinal study. *Global Change Biology*, 18, 1468-1479. doi:10.1111/j.1365-2486.2011.02615.x.
- *Gong, S., Zhang, T., Guo, R., Cao, H., Shi, L., Guo, J., & Sun, W. (2015). Response of soil enzyme activity to warming and nitrogen addition in a meadow steppe. *Soil Research*, 53, 242-252. doi:10.1071/sr14140.
- *Gutknecht, J. L., Henry, H. A., & Balsler, T. C. (2010). Inter-annual variation in soil extra-cellular enzyme activity in response to simulated global change and fire disturbance. *Pedobiologia*, 53, 283-293. doi:10.1016/j.pedobi.2010.02.001.
- Hanson, P. J., Riggs, J. S., Nettles, W. R., Phillips, J. R., Krassovski, M. B., Hook, L. A., . . . Ricciuto, D. M. (2017). Attaining whole-ecosystem warming using air and deep-soil heating methods with an elevated CO₂ atmosphere. *Biogeosciences*, 14, 861. doi:10.5194/bg-14-861-2017.
- Hedges, L. V., Gurevitch, J., & Curtis, P. S. (1999). The meta-analysis of response ratios in experimental ecology. *Ecology*, 80, 1150-1156. doi:10.1890/0012-9658(1999)080[1150:TMAORR]2.0.CO;2.
- Hobbie, S. E. (2008). Nitrogen effects on decomposition: A five-year experiment in eight temperate sites. *Ecology*, 89, 2633-2644. doi:10.1890/07-1119.1.
- *Hou, R., Zhu, O., Maxim, D., Wilson, G., & Kuzyakov, Y. (2016). Lasting effect of soil warming on organic matter decomposition depends on tillage practices. *Soil Biology and Biochemistry*, 95, 243-249. doi:10.1016/j.soilbio.2015.12.008.
- Janssens, I., Dieleman, W., Luysaert, S., Subke, J. A., Reichstein, M., Ceulemans, R., . . . Matteucci, G. (2010). Reduction of forest soil respiration in response to nitrogen deposition. *Nature Geoscience*, 3, 315-322. doi:10.1038/ngeo844.

- *Jassey, V. E. J., Chiapusio, G., Gilbert, D., Toussaint, M. L., & Binet, P. (2012). Phenoloxidase and peroxidase activities in Sphagnum-dominated peatland in a warming climate. *Soil Biology and Biochemistry*, 46, 49-52. doi:10.1016/j.soilbio.2011.11.011.
- Jian, S., Li, J., Chen, J., Wang, G., Mayes, M. A., Dzantor, K. E., . . . Luo, Y. (2016). Soil extracellular enzyme activities, soil carbon and nitrogen storage under nitrogen fertilization: A meta-analysis. *Soil Biology and Biochemistry*, 101, 32-43. doi:10.1016/j.soilbio.2016.07.003.
- *Jiang, R., Yu, Y., Tang, Y. R., Hua, Z. J., Xian, J. R., & Yang, Z. B. (2018) Effects of warming and biochar addition on soil enzyme activities in farmland. *Journal of Sichuan Agricultural University*, 36, 72-85. Doi:10.16036/j.issn.1000-2650.2018.01.011. (In Chinese with English abstract)
- *Jing, X., Wang, Y., Chung, H., Mi, Z., Wang, S., Zeng, H., & He, J. S. (2014). No temperature acclimation of soil extracellular enzymes to experimental warming in an alpine grassland ecosystem on the Tibetan Plateau. *Biogeochemistry*, 117, 39-54. doi:10.1007/s10533-013-9844-2.
- *Kane, E. S., Mazzoleni, L. R., Kratz, C. J., Hribljan, J. A., Johnson, C. P., Pypker, T. G., & Chimner, R. (2014). Peat porewater dissolved organic carbon concentration and lability increase with warming: a field temperature manipulation experiment in a poor-fen. *Biogeochemistry*, 119, 161-178. doi:10.1007/s10533-014-9955-4.
- *Kardol, P., Cregger, M. A., Company, C. E., & Classen, A. T. (2010). Soil ecosystem functioning under climate change: plant species and community effects. *Ecology*, 91, 767-781. doi:10.1890/09-0135.1.
- Karhu, K., Auffret, M. D., Dungait, J. a. J., Hopkins, D. W., Prosser, J. I., Singh, B. K., . . . Hartley, I. P. (2014). Temperature sensitivity of soil respiration rates enhanced by microbial community response. *Nature*, 513, 81-84. doi:10.1038/Nature13604.
- Kinnunen, A., Maijala, P., Jarvinen, P., & Hatakka, A. (2017). Improved efficiency in screening for lignin-modifying peroxidases and laccases of basidiomycetes. *Current Biotechnology*, 6, 105-115. doi:10.2174/2211550105666160330205138.
- *Li, N., Wang, G., Gao, Y., & Wang, J. (2011). Warming effects on plant growth, soil nutrients, microbial biomass and soil enzymes activities of two alpine meadows in Tibetan plateau. *Polish Journal of Ecology*, 59, 25-35. <https://www.infona.pl/resource/bwmeta1.element.baztech-article-BGPK-3205-2423>.
- Lin, J., Zhu, B., & Cheng, W. (2015). Decadally cycling soil carbon is more sensitive to warming than faster-cycling soil carbon. *Global Change Biology*, 21, 4602-4612. doi:10.1111/gcb.13071.

- *Liu, L., Zhu, X., Sun, G., Luo, P., & Wang, B. (2011). Effects of simulated warming and fertilization on activities of soil enzymes in alpine meadow. *Pratacultural Science*, 28, 1405-1410. doi:1001-0629(2011).28:8<1405:mznwys>2.0.tx;2-m. (In Chinese with English abstract)
- Lu, M., Zhou, X., Yang, Q., Li, H., Luo, Y., Fang, C., . . . Li, B. (2013) Responses of ecosystem carbon cycle to experimental warming: A meta-analysis. *Ecology*, 94, 726-738. doi:10.1890/12-0279.1.
- Luo, Y., Ahlström, A., Allison, S. D., Batjes, N. H., Brovkin, V., Carvalhais, N., . . . Finzi, A. (2016). Toward more realistic projections of soil carbon dynamics by Earth system models. *Global Biogeochemical Cycles*, 30, 40-56. doi:10.1002/2015GB005239.
- Luo, Y., Chen, J., Chen, Y., & Feng, W. (2017). Data-driven microbial modeling for soil carbon decomposition and stabilization. In: *EGU General Assembly Conference Abstracts*. <https://meetingorganizer.copernicus.org/EGU2017/EGU2017-4548.pdf>.
- Luo, Y., Wan, S., Hui, D., & Wallace, L. L. (2001). Acclimatization of soil respiration to warming in a tall grass prairie. *Nature*, 413, 622-625. doi:10.1038/35098065.
- *Machmuller, M. B., Mohan, J. E., Minucci, J. M., Phillips, C. A., & Wurzbürger, N. (2016). Season, but not experimental warming, affects the activity and temperature sensitivity of extracellular enzymes. *Biogeochemistry*, 1-11. doi:10.1007/s10533-016-0277-6.
- Manzoni, S., Taylor, P., Richter, A., Porporato, A., & Gren, G. I. (2012). Environmental and stoichiometric controls on microbial carbon-use efficiency in soils. *New Phytologist*, 196, 79. doi:10.1111/j.1469-8137.2012.04225.x.
- *Mcdaniel, M. D., Kaye, J. P., & Kaye, M. W. (2013). Increased temperature and precipitation had limited effects on soil extracellular enzyme activities in a post-harvest forest. *Soil Biology and Biochemistry*, 56, 90-98. doi:10.1016/j.soilbio.2012.02.026.
- Melillo, J. M., Butler, S., Johnson, J., Mohan, J., Steudler, P., Lux, H., . . . Scott, L. (2011). Soil warming, carbon–nitrogen interactions, and forest carbon budgets. *Proceedings of the National Academy of Sciences*, 108, 9508-9512. doi:10.1073/pnas.1018189108.
- Melillo, J. M., Frey, S. D., Deangelis, K. M., Werner, W. J., Bernard, M. J., Bowles, F., . . . Grandy, A. (2017). Long-term pattern and magnitude of soil carbon feedback to the climate system in a warming world. *science*, 358, 101-105. doi:10.1126/science.aan2874.
- Metcalfe, D. B. (2017). Microbial change in warming soils. *science*, 358, 41-42. doi:10.1126/science.aap7325.

- Moorhead, D. L., Sinsabaugh, R. L., Hill, B. H., & Weintraub, M. N. (2016). Vector analysis of ecoenzyme activities reveal constraints on coupled C, N and P dynamics. *Soil Biology and Biochemistry*, 93, 1-7. doi:10.1016/j.soilbio.2015.10.019.
- *Nie, M., Pendall, E., Bell, C., Gasch, C. K., Raut, S., Tamang, S., & Wallenstein, M. D. (2013). Positive climate feedbacks of soil microbial communities in a semi-arid grassland. *Ecology Letters*, 16, 234-241. doi:10.1111/Ele.12034.
- *Nie, M., Pendall, E., Bell, C., & Wallenstein, M. D. (2014). Soil aggregate size distribution mediates microbial climate change feedbacks. *Soil Biology and Biochemistry*, 68, 357-365. doi:10.1016/j.soilbio.2013.10.012.
- O'Neill, B. C., Oppenheimer, M., Warren, R., Hallegatte, S., Kopp, R. E., Pörtner, H. O., . . . Licker, R. (2017). IPCC reasons for concern regarding climate change risks. *Nature Climate Change*, 7, 28-37. doi:10.1038/nclimate3179.
- *Pan, X., Lin, B., & Liu, Q. (2008). Effects of elevated temperature on soil organic carbon and soil respiration under subalpine coniferous forest in western Sichuan Province, China. *Chinese Journal of Applied Ecology*, 19, 1637-1643. <http://www.cjae.net/CN/Y2008/V19/I08/1637>. (In Chinese with English abstract)
- *Papanikolaou, N., Britton, A. J., Helliwell, R. C., & Johnson, D. (2010). Nitrogen deposition, vegetation burning and climate warming act independently on microbial community structure and enzyme activity associated with decomposing litter in low-alpine heath. *Global Change Biology*, 16, 3120-3132. doi:10.1111/j.1365-2486.2010.02196.x.
- Paustian, K., Lehmann, J., Ogle, S., Reay, D., Robertson, G. P., & Smith, P. (2016). Climate-smart soils. *Nature*, 532, 49-57. doi:10.1038/nature17174.
- Peñuelas, J., Ciais, P., Canadell, J. G., Janssens, I. A., Fernández-Martínez, M., Carnicer, J., . . . Sardans, J. (2017). Shifting from a fertilization-dominated to a warming-dominated period. *Nature Ecology & Evolution*, 1, 1438. doi:10.1038/s41559-017-0274-8.
- *Pold, G., Grandy, A. S., Melillo, J. M., & Deangelis, K. M. (2017). Changes in substrate availability drive carbon cycle response to chronic warming. *Soil Biology and Biochemistry*, 110, 68-78. doi:10.1016/j.soilbio.2017.03.002.
- *Pold, G., Melillo, J. M., & Deangelis, K. M. (2015). Two decades of warming increases diversity of a potentially lignolytic bacterial community. *Frontiers in Microbiology*, 6. doi:10.3389/fmicb.2015.00480.
- *Qiao, M., Xiao, J., Yin, H., Pu, X., Yue, B., & Liu, Q. (2014). Analysis of the phenolic compounds in root exudates produced by a subalpine coniferous species as responses to experimental warming

and nitrogen fertilisation. *Chemistry and Ecology*, 30, 555-565.
doi:10.1080/02757540.2013.868891.

*Ren, X. W., Tang, J. Y., Liu, J. C., Huai-Jiang, H. E., Dong, D., & Cheng, Y. X. (2014). Effects of elevated CO₂ and temperature on soil enzymes of seedlings under different nitrogen concentrations. *Journal of Beijing Forestry University*, 36, 44-53. doi:10.13332/j.cnki.jbfu.2014.05.016. (In Chinese with English abstract)

*Romero-Olivares, A. L., Allison, S. D., & Treseder, K. K. (2017). Decomposition of recalcitrant carbon under experimental warming in boreal forest. *Plos One*, 12, e0179674.
doi:10.1371/journal.pone.0179674.

Rousk, J., Smith, A. R., & Jones, D. L. (2013). Investigating the long-term legacy of drought and warming on the soil microbial community across five European shrubland ecosystems. *Global Change Biology*, 19, 3872-3884. doi:10.1111/gcb.12338.

Rustad, L. E., Campbell, J. L., Marion, G. M., Norby, R. J., Mitchell, M. J., Hartley, A. E., . . . Gcte, N. (2001). A meta-analysis of the response of soil respiration, net nitrogen mineralization, and aboveground plant growth to experimental ecosystem warming. *Oecologia*, 126, 543-562.
doi:10.1007/s004420000544.

*Sang, L. (2017) Effects of warming and straw application on soil respiration and enzyme activities in a cropland. *Nanjing University of Information Science & Technology*.
<http://cdmd.cnki.com.cn/Article/CDMD-10300-1017295075.htm>. (In Chinese with English abstract)

*Sardans, J., Penuelas, J., & Estiarte, M. (2008). Changes in soil enzymes related to C and N cycle and in soil C and N content under prolonged warming and drought in a Mediterranean shrubland. *Applied Soil Ecology*, 39, 223-235. doi:10.1016/j.apsoil.2007.12.011.

Schindlbacher, A., Rodler, A., Kuffner, M., Kitzler, B., Sessitsch, A., & Zechmeister-Boltenstern, S. (2011). Experimental warming effects on the microbial community of a temperate mountain forest soil. *Soil Biology and Biochemistry*, 43, 1417-1425. doi:10.1016/j.soilbio.2011.03.005.

*Schindlbacher, A., Schneckner, J., Takriti, M., Borken, W., & Wanek, W. (2015). Microbial physiology and soil CO₂ efflux after 9 years of soil warming in a temperate forest—no indications for thermal adaptations. *Global Change Biology*, 21, 4265-4277. doi:10.1111/gcb.12996.

Schindlbacher, A., Wunderlich, S., Borken, W., Kitzler, B., Zechmeister-Boltenstern, S., & Jandl, R. (2012). Soil respiration under climate change: prolonged summer drought offsets soil warming effects. *Global Change Biology*, 18, 2270-2279.
doi:10.1111/j.1365-2486.2012.02696.x.

- *Schuerings, J., Jentsch, A., Hammerl, V., Lenz, K., Henry, H. A. L., Malyshev, A. V., & Kreyling, J. (2014). Increased winter soil temperature variability enhances nitrogen cycling and soil biotic activity in temperate heathland and grassland mesocosms. *Biogeosciences*, 11, 7051-7060. doi:10.5194/bg-11-7051-2014.
- Sinsabaugh, R. L. (2010). Phenol oxidase, peroxidase and organic matter dynamics of soil. *Soil Biology and Biochemistry*, 42, 391-404. doi:10.1016/j.soilbio.2009.10.014.
- Sinsabaugh, R. L., Lauber, C. L., Weintraub, M. N., Ahmed, B., Allison, S. D., Crenshaw, C., . . . Gallo, M. E. (2008). Stoichiometry of soil enzyme activity at global scale. *Ecology Letters*, 11, 1252-1264. doi:10.1111/j.1461-0248.2008.01245.x.
- *Sistla, S. A. & Schimel, J. P. (2013). Seasonal patterns of microbial extracellular enzyme activities in an arctic tundra soil: Identifying direct and indirect effects of long-term summer warming. *Soil Biology and Biochemistry*, 66, 119-129. doi:10.1016/j.soilbio.2013.07.003.
- *Sorensen, P. O., Finzi, A. C., Giasson, M. A., Reinman, A. B., Sanders-Demot, R., & Temple, P. H. (2018). Winter soil freeze-thaw cycles lead to reductions in soil microbial biomass and activity not compensated for by soil warming. *Soil Biology and Biochemistry*, 116, 39-47. doi:10.1016/j.soilbio.2017.09.026.
- *Souza, R. C., Solly, E. F., Dawes, M. A., Graf, F., Hagedorn, F., Egli, S., . . . Peter, M. (2017). Responses of soil extracellular enzyme activities to experimental warming and CO₂ enrichment at the alpine treeline. *Plant & Soil*, 1-11. doi:10.1007/s11104-017-3235-8.
- *Steinauer, K., Tilman, D., Wragg, P. D., Cesarz, S., Cowles, J. M., Pritsch, K., . . . Eisenhauer, N. (2015). Plant diversity effects on soil microbial functions and enzymes are stronger than warming in a grassland experiment. *Ecology*, 96, 99-112. doi:10.1890/14-0088.1.
- *Steinweg, J. M., Dukes, J. S., Paul, E. A., & Wallenstein, M. D. (2013). Microbial responses to multi-factor climate change: effects on soil enzymes. *Frontiers in Microbiology*, 4. doi:10.3389/fmicb.2013.00146.
- Stone, M. M., Weiss, M. S., Goodale, C. L., Adams, M. B., Fernandez, I. J., German, D. P., & Allison, S. D. (2012). Temperature sensitivity of soil enzyme kinetics under N-fertilization in two temperate forests. *Global Change Biology*, 18, 1173-1184. doi:10.1111/j.1365-2486.2011.02545.x.
- *Sun, H., Wu, X. C., Qin, J. H., & Yang, W. (2007) Response of soil catalase activities to temperature and CO₂ in subalpine forest in the western Sichuan. *Chinese Journal of Soil Science*, 38, 891-895. doi: 10.5846/stxb201109251406. (In Chinese with English abstract)

- Suseela, V., Tharayil, N., Xing, B. S., & Dukes, J. S. (2014). Warming alters potential enzyme activity but precipitation regulates chemical transformations in grass litter exposed to simulated climatic changes. *Soil Biology and Biochemistry*, 75, 102-112. doi:10.1016/j.soilbio.2014.03.022.
- Talbot, J. M., Yelle, D. J., Nowick, J., & Treseder, K. K. (2012). Litter decay rates are determined by lignin chemistry. *Biogeochemistry*, 108, 279-295. doi:10.1007/s10533-011-9599-6.
- Terrer, C., Vicca, S., Hungate, B. A., Phillips, R. P., & Prentice, I. C. (2016). Mycorrhizal association as a primary control of the CO₂ fertilization effect. *Science*, 353, 72-74. doi:10.1126/science.aaf4610.
- Tucker, C. L., Bell, J., Pendall, E., & Ogle, K. (2013). Does declining carbon-use efficiency explain thermal acclimation of soil respiration with warming? *Global Change Biology*, 19, 252-263. doi:10.1111/gcb.12036.
- Van Gestel, N., Shi, Z., van Groenigen, K. J., Osenberg, C.W., Andresen, L. C., Dukes, J. S., . . . Hungate, B. A. (2018). Predicting soil carbon loss with warming. *Nature*, 554, E4. doi:10.1038/nature25745.
- Van Groenigen, K. J., Qi, X., Osenberg, C. W., Luo, Y., & Hungate, B. A. (2014). Faster decomposition under increased atmospheric CO₂ limits soil carbon storage. *Science*, 344, 508-509. doi:10.1126/science.1249534.
- Van Groenigen, K. J., Osenberg, C. W., Terrer, C., Carrillo, Y., Dijkstra, F., Heath, J., . . . Hungate, B. A. (2017). Faster turnover of new soil carbon inputs under increased atmospheric CO₂. *Global Change Biology*, 23, 4420-4429. doi:10.1111/gcb.13752.
- Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*, 36, 1-48.
- Waldrop, M. P., Zak, D. R., Sinsabaugh, R. L., Gallo, M., & Lauber, C. (2004). Nitrogen deposition modifies soil carbon storage through changes in microbial enzymatic activity. *Ecological Applications*, 14, 1172-1177. doi:10.1890/03-5120.
- *Wang, H., He, Z., Lu, Z., Zhou, J., Van Nostrand, J. D., Xu, X., & Zhang, Z. (2012). Genetic linkage of soil carbon pools and microbial functions in subtropical freshwater wetlands in response to experimental warming. *Applied and Environmental Microbiology*, 78, 7652-7661. doi:10.1128/aem.01602-12.
- *Wang, X., Zhou, Y., Wang, X., Jiang, X., & Han, S. (2014a). Responses of soil enzymes in activity and soil microbes in biomass to warming in tundra ecosystems on Changbai mountains. *Acta Pedologica Sinica*, 51, 166-175. doi:10.11766/trxb201303120112. (In Chinese with English abstract)

- *Wang, X., Dong, S., Gao, Q., Zhou, H., Liu, S., Su, X., & Li, Y. (2014b). Effects of short-term and long-term warming on soil nutrients, microbial biomass and enzyme activities in an alpine meadow on the Qinghai-Tibet Plateau of China. *Soil Biology and Biochemistry*, 76, 140-142. doi:10.1016/j.soilbio.2014.05.014.
- *Wang, Y., Liu, Y. C., & Liu, S. R. (2017) Response of soil enzyme activities to soil warming and explanation of environmental factors in warm-temperate Oak forest. *Forest Research*, 30, 117-124. Doi:10.13275/j.cnki.lykxyj.2017.01.016. (In Chinese with English abstract)
- *Weedon, J. T., Aerts, R., Kowalchuk, G. A., & Van Bodegom, P. M. (2014). No effects of experimental warming but contrasting seasonal patterns for soil peptidase and glycosidase enzymes in a sub-arctic peat bog. *Biogeochemistry*, 117, 55-66. doi:10.1007/s10533-013-9870-0.
- Weedon, J. T., Kowalchuk, G. A., Aerts, R., Van Hal, J., Van Logtestijn, R., Tas, N., . . . Van Bodegom, P. M. (2012). Summer warming accelerates sub-arctic peatland nitrogen cycling without changing enzyme pools or microbial community structure. *Global Change Biology*, 18, 138-150. doi:10.1111/j.1365-2486.2011.02548.x.
- Wieder, W. R., Bonan, G. B., & Allison, S. D. (2013). Global soil carbon projections are improved by modelling microbial processes. *Nature Climate Change*, 3, 909. doi:10.1038/nclimate1951.
- *Xu, G., Chen, J., Berninger, F., Pumpanen, J., Bai, J., Yu, L., & Duan, B. (2015). Labile, recalcitrant, microbial carbon and nitrogen and the microbial community composition at two Abies faxoniana forest elevations under elevated temperatures. *Soil Biology and Biochemistry*, 91, 1-13. doi:10.1016/j.soilbio.2015.08.016.
- *Xu, Z. F., Tang, Z., Wan, C., Xiong, P., Cao, G., & Liu, Q. (2010). Effects of simulated warming on soil enzyme activities in two subalpine coniferous forests in west Sichuan. *Chinese Journal of Applied Ecology*, 21, 2727-2733. <http://www.cjae.net/CN/Y2010/V21/I11/2727>. (In Chinese with English abstract)
- Xue, K., Yua, M.M., Sh, Z. J., Qi, Y., Den, Y., Cheng, L., . . . Bracho, R. (2016). Tundra soil carbon is vulnerable to rapid microbial decomposition under climate warming. *Nature Climate Change*, 6, 595-600. doi:10.1038/NCLIMATE2940.
- *Yang, L., Chen, Y. M., He, R. L., Deng, C. C., Liu, J. W., & Liu, Y. (2016) Responses of soil microbial community structure and function to simulated warming in alpine forest. *Chinese Journal of Applied Ecology*, 27, 2855-2863. doi:10.13287/j.1001-9332.201609.026. (In Chinese with English abstract)
- Yang, Y., Hopping, K. A., Wang, G., Chen, J., Peng, A., & Klein, J. A. (2018) Permafrost and drought regulate vulnerability of Tibetan Plateau grasslands to warming. *Ecosphere*, 9, e02233. doi:org/10.1002/ecs2.2233.

- *Yin, H., Li, Y., Xiao, J., Xu, Z., Cheng, X., & Liu, Q. (2013). Enhanced root exudation stimulates soil nitrogen transformations in a subalpine coniferous forest under experimental warming. *Global Change Biology*, 19, 2158-2167. doi:10.1111/gcb.12161.
- *Zhao, C., Zhu, L., Liang, J., Yin, H., Yin, C., Li, D., . . . Liu, Q. (2014). Effects of experimental warming and nitrogen fertilization on soil microbial communities and processes of two subalpine coniferous species in Eastern Tibetan Plateau, China. *Plant and Soil*, 382, 189-201. doi:10.1007/s11104-014-2153-2.
- *Zhao, C. Z., & Liu, Q. (2009). Growth and physiological responses of *Picea asperata* seedlings to elevated temperature and to nitrogen fertilization. *Acta Physiologiae Plantarum*, 31, 163-173. doi:10.1007/s11738-008-0217-8.
- Zhao, F., Ren, C., Shelton, S., Wang, Z., Pang, G., Chen, J., & Wang, J. (2017). Grazing intensity influence soil microbial communities and their implications for soil respiration. *Agriculture, Ecosystems & Environment*, 249, 50-56. doi:10.1016/j.agee.2017.08.007.
- *Zhou, J. Z., Xue, K., Xie, J. P., Deng, Y., Wu, L. Y., Cheng, X. H., . . . Luo, Y. Q. (2012). Microbial mediation of carbon-cycle feedbacks to climate warming. *Nature Climate Change*, 2, 106-110. doi:10.1038/Nclimate1331.
- *Zhou, X. Q., Chen, C. R., Wang, Y. F., Xu, Z. H., Han, H. Y., Li, L. H., & Wan, S. Q. (2013). Warming and increased precipitation have differential effects on soil extracellular enzyme activities in a temperate grassland. *Science of the Total Environment*, 444, 552-558. doi:10.1016/j.scitotenv.2012.12.023.

Figure captions

FIGURE 1 (a) Effects of warming on cellulase and ligninase activities indicated with the mean percentage of change in warming vs. control plots. Distribution of the log-transformed response ratios (lnR) of (b) cellulase and (c) ligninase activities to experimental warming. Error bars represent 95% confidence intervals. The sample size for each variable is shown in the right column of the figure. PER, peroxidase; PO, phenol oxidase; PPO, polyphenol oxidase; BG, β -1,4-Glucosidase; BX, β -1,4-Xylosidase; CBH, β -D-Cellobiosidase.

FIGURE 2 Model-averaged importance of the predictors of warming effects on soil (a) cellulase and (b) ligninase activities. The importance is based on the sum of Akaike weights derived from model selection using corrected Akaike's Information Criteria. Cutoff is set at 0.8 in order to differentiate between important and non-essential predictors. MAT, mean annual temperature; MAP, mean annual precipitation; Sample.T, sampling temperature; Time, daily warming regime (i.e., day, night, or diurnal warming); Season, annual warming regime (i.e. growing season, non-growing season or whole-year warming).

FIGURE 3 Relationships between warming-induced changes in (a) cellulase and (b) ligninase activities and warming duration. Effects of warming on (c) cellulase and (d) ligninase activities for various warming methods. The response of ligninase activity was positively correlated with warming duration ($y = 0.016x + 0.113$, $R^2 = 0.117$, $p < 0.001$, $F = 22.590$, $n = 172$). Error bars represent 95% confidence intervals. OTC, open top chamber; IH, infrared heater; GH, green house. The sample size for each variable is shown in the right column of the figure.

FIGURE 4 Relationships between the effect of warming (lnR) on soil respiration (SR) and lnR of (a) cellulase and (b) ligninase activities. (c) Model-averaged importance of the predictors of warming effects on SR. The warming response of SR was positively correlated with the warming response of ligninase activity ($y = 0.528x + 0.108$, $R^2 = 0.467$, $p < 0.001$, $F = 61.260$, $n = 72$). Model selection analysis is limited to studies that simultaneously reported the responses of ligninase, cellulase and SR. The importance is based on the sum of Akaike weights derived from model selection using corrected Akaike's Information Criteria. Cutoff is set at 0.8 to differentiate between important and non-essential predictors. MAT, mean annual temperature; MAP, mean annual precipitation; Time, daily warming regime (i.e., day, night, or diurnal warming); Season, annual warming regime (i.e. growing season, non-growing season or whole-year warming); Sample.T, sampling temperature.







